

A. E. Choque-Rivero

PhD math, Prof.

Instituto de Física y Matemáticas, Universidad Michoacana de San Nicolás de Hidalgo
 Edificio C-3, C.U., CP 58060, Morelia, Michoacán, México

abdon.choque@umich.mx  <http://orcid.org/0000-0003-0226-9612>

On the Hurwitz Stability of Hurwitz-Type Matrix Polynomials

A real scalar polynomial whose roots lie in the left half-plane of the complex plane is called a Hurwitz polynomial. This notion goes back to the works of J. C. Maxwell, E. J. Routh, and A. Hurwitz, and was later studied using techniques such as Sturm sequences, Markov parameters, and continued fractions; see, for instance, *The Theory of Matrices*, Vol. II, Chapter XV, AMS Chelsea (2000), by F. Gantmacher.

A $q \times q$ matrix polynomial $P(z)$ is called Hurwitz if $\det P(z)$ is a Hurwitz polynomial. Every matrix polynomial $\mathbf{f}_n$ can be written in the form $\mathbf{f}_n(z) = \mathbf{h}_n(z^2) + z \mathbf{g}_n(z^2)$. The matrix polynomial $\mathbf{f}_{2m}$ is said to be of Hurwitz type if the expression $\mathbf{g}_{2m}(z) \mathbf{h}_{2m}^{-1}(z)$ admits a representation as a finite continued fraction with positive definite matrix coefficients. Similarly, the odd-degree matrix polynomial $\mathbf{f}_{2m+1}$ is of Hurwitz type if $\frac{1}{z} \mathbf{h}_{2m+1}(z) \mathbf{g}_{2m+1}^{-1}(z)$ has the same property. The concept of Hurwitz-type matrix polynomials was introduced in *On matrix Hurwitz type polynomials and their interrelations to Stieltjes positive definite sequences and orthogonal matrix polynomials*, Linear Algebra Appl. 476 (2015), by A. E. Choque Rivero.

In the present work, we derive an explicit form of the Bezoutian associated with Hurwitz-type matrix polynomials. The fact that Hurwitz-type matrix polynomials are Hurwitz matrix polynomials was suggested and partially proved using Bezoutians in *On generalization of classical Hurwitz stability criteria for matrix polynomials*, J. Comput. Appl. Math. 383 (2021), by X. Zhan and A. Dyachenko. In contrast to that work, we employ the decomposition of the Bezoutian form introduced in *Some Questions in the Theory of Moments*, Translations of Mathematical Monographs 2, AMS, 1962, by N. I. Akhiezer and M. G. Krein, for scalar polynomials.

Additionally, we propose a method to enlarge the class of Hurwitz-type matrix polynomials by adding to a given polynomial a matrix polynomial that is not of Hurwitz type, so that the resulting polynomial becomes of Hurwitz type.

Keywords: Hurwitz matrix polynomial; orthogonal matrix polynomial; Bezoutian; Hurwitz-Type Matrix Polynomial.

2020 Mathematics Subject Classification: 34D20; 34D20; 33C45; 47A56; 15A24; 30E05

1. Introduction

A scalar polynomial with all its roots in the left half of the complex plane is called a Hurwitz polynomial [47, 34, 16, 27, 33, 36, 8, 35]. The Hurwitz stability of such polynomials is also referred to as Hurwitzness—a term we adopt in this work. Let n and q be natural numbers. In the present work, we consider the properties and Hurwitzness of a subset of Lambda matrices [43, page 95], [28, page 228], [17], defined by

$$\mathbf{f}_n(z) := A_0 z^n + A_1 z^{n-1} + \dots + A_n, \quad (1)$$

where A_j is a complex $q \times q$ matrix and z is a complex variable. As in [39], we refer to a matrix of the form (1) as a matrix polynomial, which is called a Hurwitz matrix polynomial if $\det \mathbf{f}_n$ is a Hurwitz polynomial.

The matrix polynomial (1) has degree n if A_0 is nonzero. Throughout this paper, we assume that $\det A_0 \neq 0$. Clearly, (1) can be represented as

$$\mathbf{f}_n(z) = \mathbf{h}_n(z^2) + z \mathbf{g}_n(z^2), \quad (2)$$

where

$$\mathbf{h}_n(z) := \begin{cases} A_0 z^m + A_2 z^{m-1} + \dots + A_{2m}, & n = 2m, \\ A_1 z^m + A_3 z^{m-1} + \dots + A_{2m+1}, & n = 2m + 1, \end{cases} \quad (3)$$

and

$$\mathbf{g}_n(z) := \begin{cases} A_1 z^{m-1} + A_3 z^{m-2} + \dots + A_{2m-1}, & n = 2m, \\ A_0 z^m + A_2 z^{m-1} + \dots + A_{2m}, & n = 2m + 1. \end{cases} \quad (4)$$

We will refer to (3) as the $\mathbf{h}_n$ part and to (4) as the $\mathbf{g}_n$ part of the matrix polynomial $\mathbf{f}_n$.

Let us introduce the following notation. The symbol $\mathbb{C}^{q \times q}$ represents the set of all complex $q \times q$ matrices. For the null matrix that belongs to $\mathbb{C}^{p \times q}$, we will write $0_{p \times q}$. We denote by 0_q and I_q the null and identity matrices in $\mathbb{C}^{q \times q}$, respectively. When the sizes of the null and identity matrices are clear from the context, we will omit their indices.

In the following definition, we recall the concept of a Hurwitz-type matrix polynomial (HTM polynomial) introduced in [6]. In [6], HTM polynomials were referred to as matrix Hurwitz-type polynomials.

For $A, B \in \mathbb{C}^{q \times q}$ with B invertible, set

$$\frac{A}{B} := AB^{-1}. \quad (5)$$

Definition 1. The $q \times q$ matrix polynomial $\mathbf{f}_n$ in (1) is called a Hurwitz-type matrix polynomial if there exist two sequences of positive definite matrices, $(\mathbf{c}_k)_{k=0}^{m-1}$ and $(\mathbf{d}_k)_{k=0}^{m-1}$ (resp. $(\mathbf{c}_k)_{k=0}^m$ and $(\mathbf{d}_k)_{k=0}^{m-1}$) such that, for $n = 2m$,

$$\frac{\mathbf{g}_n(z)}{\mathbf{h}_n(z)} = \frac{I_q}{z\mathbf{c}_0 + \frac{I_q}{\mathbf{d}_0 + \frac{I_q}{\ddots + \mathbf{d}_{m-2} + \frac{I_q}{+z\mathbf{c}_{m-1} + \mathbf{d}_{m-1}^{-1}}}}}, \quad (6)$$

for all $z \in \mathbb{C}$ with $\det \mathbf{h}_n(z) \neq 0$ and, for $n = 2m + 1$,

$$\frac{\mathbf{h}_n(z)}{z\mathbf{g}_n(z)} = \frac{I_q}{z\mathbf{c}_0 + \frac{I_q}{\mathbf{d}_0 + \frac{I_q}{\ddots + z\mathbf{c}_{m-1} + \frac{I_q}{\mathbf{d}_{m-1} + z^{-1}\mathbf{c}_m^{-1}}}}}, \quad (7)$$

for all $z \in \mathbb{C} \setminus \{0\}$ with $\det \mathbf{g}_n(z) \neq 0$.

The definition of HTM polynomials in the form of a finite continued fraction is inspired by [29, Chapter XV, Theorem 15].

Main Results. Below are the main results of the present work:

- (a) We compute explicitly the Bezoutian of the HTM polynomial (1); see Theorem 1 and Corollary 2. From this, an explicit proof of the Hurwitz property of HTM polynomials readily follows; see Theorem 2.
- (b) For a given matrix polynomial P_n of degree n that is not a Hurwitz-type matrix (HTM) polynomial, we construct a polynomial $\mathbf{f}_{2n}(z) = P_n(z^2) + zQ_{n-1}(z^2)$ of degree $2n$, with a suitably chosen Q_{n-1} , such that $\mathbf{f}_{2n}$ becomes an HTM polynomial. Then, by applying the result from [21], we conclude that P_n itself is a Hurwitz matrix polynomial; see Theorem 4.

Relation between (a) and (b). The explicit computation of the Bezoutian in part (a) provides the structural foundation for the Hurwitz property of HTM polynomials. This result is then exploited in part (b), where a general matrix polynomial is embedded into an HTM polynomial of higher degree. The Hurwitz property established in part (a), together with the construction in (b), allows us to transfer stability from the HTM setting back to the original polynomial P_n .

The Hurwitz property of matrix polynomials via Bezoutians was studied in [30, 41, 42]. In [49], the Hurwitz property of HTM polynomials was also considered. However, the proof of [49, Theorem 3.5], which serves as the basis for the analysis in that work, is not fully detailed. More precisely, several steps in the proof are

left implicit; see Remark 14. In particular, the argument is developed only for the case $n = 2m$, while the case $n = 2m + 1$ is described as “similar,” without explicit justification.

A distinctive feature of the odd-degree case $n = 2m + 1$ is that the associated block Hankel matrices $H_{1,m}$ and $H_{2,m-1}$ have different dimensions. In contrast to the even case, this asymmetry prevents a direct application of the standard construction. To overcome this difficulty, we introduce a modified polynomial $\tilde{\mathbf{g}}_{2m+1}$ (see (57)) of degree $m - 1$, which replaces the original polynomial $\mathbf{g}_{2m+1}$ (of degree m) in the construction of the symmetrizer. This approach leads to a consistent formulation of the matrix $\mathcal{W}_{2m+1}^{(0)}$ (see (76) and Remark 5) and provides an explicit treatment of the odd-degree case, which is typically addressed only heuristically in the literature. This aspect is not discussed in [49].

Additionally, in contrast to [49], we use representations of the matrix polynomial $\mathbf{f}_n$ in terms of orthogonal polynomials on $[0, \infty)$ of the first kind and polynomials of the second kind, as in (2) and (18), for example, to compute explicitly the Bezoutian of the quadruple $(\mathbf{f}_n^*(\bar{x}), \mathbf{f}_n(-x), \mathbf{f}_n^*(-\bar{x}), \mathbf{f}_n(x))$.

The asterisk denotes the Hermitian adjoint of a matrix, while the overbar represents complex conjugation. In the scalar case, the use of the $\mathbf{h}_n$ part and $\mathbf{g}_n$ part of the polynomial $\mathbf{f}_n$ for studying the corresponding Bezoutian was employed by Krein and Naimark in [38] and by Krein and Akhiezer in [1].

The importance of HTM polynomials as a subset of Hurwitz matrix polynomials is that they help study the asymptotic stability of the differential equation

$$I_q y^{(n)}(t) + A_1 y^{(n-1)}(t) + \dots + A_{n-1} y'(t) + A_n y(t) = 0, \quad t \geq 0$$

with $y(t) \in \mathbb{R}^n$; $I_q, A_1, \dots, A_n \in \mathbb{C}^{q \times q}$; see [44, 48, 46, 31] and [37].

Our motivation is to generalize the results regarding robust stability [7] and finite-time stabilization with bounded controls [13, 14, 9, 11] for the matrix case within the framework of HTM polynomials.

Our work is organized as follows. In Section 2, we review previous results on HTM polynomials and their associated Markov parameters. We also recall Stieltjes orthogonal matrix polynomials, orthogonal matrix polynomials, the interrelations between HTM polynomials and Markov parameters, as well as the Bezoutian and the inertia of matrix polynomials. In Section 3, we explicitly compute the Bezoutian of HTM polynomials. In Section 4, we study the Hurwitzness of HTM polynomials. In Section 5, we present a result and an algorithm for constructing an HTM polynomial from a given matrix polynomial that is not of HTM type. We also explain why the original matrix polynomial is a Hurwitz matrix polynomial. In Section 6, we clarify certain steps left implicit in Theorem 3.5 of [49], concerning a “matrix analogue of the stability criterion via Markov parameters,” as stated on page 10 of [49].

This work is a direct continuation of the author’s research in [5] and [6], which focus on the nondegenerate truncated matricial Stieltjes problem and HTM polynomials, respectively.

2. Preliminaries

In this section, we introduce the notation and preliminary results that will be used throughout the paper.

HTM polynomials and associated Markov parameters

The following quantities will be used to describe the domains of validity of the Laurent expansions appearing below.

If $n = 2m$, we define

$$\mathcal{N}_{\mathbf{h}_n} := \{z \in \mathbb{C} : \det \mathbf{h}_n(z) = 0\}, \quad \eta_n := \max\{|z| : z \in \mathcal{N}_{\mathbf{h}_n}\}, \quad (8)$$

whereas, for $n = 2m + 1$, we define

$$\mathcal{N}_{\mathbf{g}_n} := \{z \in \mathbb{C} : \det \mathbf{g}_n(z) = 0\}, \quad \rho_n := \max\{|z| : z \in \mathcal{N}_{\mathbf{g}_n}\}. \quad (9)$$

The next remark recalls the Laurent expansions of $\mathbf{g}_n(z)\mathbf{h}_n(z)^{-1}$ and $\mathbf{h}_n(z)(z\mathbf{g}_n(z))^{-1}$, which connect the matrix polynomial (1) with the associated Markov parameters s_j .

Remark 1. Let $\mathbf{h}_n$ and $\mathbf{g}_n$ be defined as in (3) and (4), respectively, and let $\mathbf{f}_n$ be given by (2).

For $|z| > \eta_n$ (resp. $|z| > \rho_n$), the Laurent expansion in negative powers of z of $\mathbf{g}_n(z)\mathbf{h}_n(z)^{-1}$ (resp. $\mathbf{h}_n(z)(z\mathbf{g}_n(z))^{-1}$) takes the form

$$\frac{\mathbf{g}_n(z)}{\mathbf{h}_n(z)} = \frac{s_0}{z} - \frac{s_1}{z^2} + \cdots + (-1)^n \frac{s_n}{z^{n+1}} + \cdots, \quad n = 2m, \quad (10)$$

$$\frac{\mathbf{h}_n(z)}{z\mathbf{g}_n(z)} = \frac{s_0}{z} - \frac{s_1}{z^2} + \cdots + (-1)^n \frac{s_n}{z^{n+1}} + \cdots, \quad n = 2m + 1. \quad (11)$$

For $j \geq 0$ and by using the Markov parameters, we construct block Hankel matrices:

$$H_{1,j} := \begin{pmatrix} s_0 & s_1 & \cdots & s_j \\ s_1 & s_2 & \cdots & s_{j+1} \\ \vdots & \vdots & \ddots & \vdots \\ s_j & s_{j+1} & \cdots & s_{2j} \end{pmatrix}, \quad H_{2,j} := \begin{pmatrix} s_1 & s_2 & \cdots & s_{j+1} \\ s_2 & s_3 & \cdots & s_{j+2} \\ \vdots & \vdots & \ddots & \vdots \\ s_{j+1} & s_{j+2} & \cdots & s_{2j+1} \end{pmatrix}. \quad (12)$$

The rational matrix functions in (6) and (7) are special solutions to the truncated Stieltjes matrix moment (TSMM) problem; see proof of [6, Theorem 7.9]. This implies that $\frac{\mathbf{g}_n(-z)}{\mathbf{h}_n(-z)}$ and $\frac{\mathbf{h}_n(-z)}{(-z)\mathbf{g}_n(-z)}$ are Stieltjes transforms of a certain positive measure σ on $[0, +\infty)$.

The TSMM problem is defined as follows: Let $(s_j)_{j=0}^n$ be a finite sequence of $q \times q$ complex Hermitian matrices. The TSMM problem consists of finding the set $\mathcal{M}_n$ of all nonnegative Hermitian $q \times q$ measures σ defined on the Borel σ -algebra $\mathfrak{B} \cap [0, +\infty)$ such that

$$s_j = \int_{[0, +\infty)} t^j \sigma(dt), \quad 0 \leq j \leq n-1, \quad s_n = \int_{[0, +\infty)} t^n \sigma(dt) + M,$$

where M is a $q \times q$ complex valued positive semi-definite matrix.

For each $\sigma \in \mathcal{M}_n$, one associates a matrix-valued function

$$s(z) = \int_{[0,+\infty)} \frac{\sigma(dt)}{t-z}, \quad (13)$$

which is defined and holomorphic in $\mathbb{C} \setminus [0, +\infty)$.

The function s is associated with the TSMM problem and is called the Stieltjes transform of the positive measure σ ; see [18, 19, 20, 22, 23, 15, 24, 25, 26].

In 1981, Yu. Dyukarev proved that the TSMM problem in the case of an odd (resp. even) number of moments has a solution if and only if $H_{1,m}$ and $H_{2,m-1}$ (resp. $H_{1,m}$ and $H_{2,m}$) are both positive semidefinite matrices.

Definition 2. *The sequence $(s_j)_{j=0}^{2m}$ (resp. $(s_j)_{j=0}^{2m+1}$) is called a Stieltjes positive sequence if both corresponding matrices $H_{1,m}$ and $H_{2,m-1}$ (resp. $H_{1,m}$ and $H_{2,m}$) are positive definite. In the case where $(s_j)_{j=0}^{2m}$ (resp. $(s_j)_{j=0}^{2m+1}$) is a positive Stieltjes sequence, the corresponding TSMM problem is called a TSMM problem in the nondegenerate case.*

In [18], Yu. Dyukarev found the complete set of solutions to the TSMM problem in the case that $(s_j)_{j=0}^{2m}$ and $(s_j)_{j=0}^{2m+1}$ are Stieltjes positive sequences, respectively.

Stieltjes orthogonal matrix polynomials

We now reproduce from [5] a number of auxiliary matrices, as well as two families of orthogonal matrix polynomials and their polynomials of the second kind. Let $R_j : \mathbb{C} \rightarrow \mathbb{C}^{(j+1)q \times (j+1)q}$ be given by

$$R_j(z) := (I_{(j+1)q} - zT_j)^{-1}, \quad j \geq 0,$$

with

$$T_0 := 0_q, \quad T_j := \begin{pmatrix} 0_{q \times jq} & 0_q \\ I_{jq} & 0_{jq \times q} \end{pmatrix}, \quad j \geq 1.$$

Observe that for each $j \in \mathbb{N}_0$, the matrix-valued function R_j can be represented via

$$R_j(z) = \begin{pmatrix} I_q & 0_q & 0_q & \cdots & 0_q & 0_q \\ zI_q & I_q & 0_q & \cdots & 0_q & 0_q \\ z^2I_q & zI_q & I_q & \cdots & 0_q & 0_q \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ z^jI_q & z^{j-1}I_q & z^{j-2}I_q & \cdots & zI_q & I_q \end{pmatrix}. \quad (14)$$

Let

$$v_0 := I_q, \quad v_j := \begin{pmatrix} I_q \\ 0_{jq \times q} \end{pmatrix} \text{ for all } j \in \mathbb{N}. \quad (15)$$

Furthermore, let

$$Y_{1,j} := y_{[j,2j-1]}, \quad 1 \leq j \leq n, \quad \text{and} \quad Y_{2,j} := y_{[j+1,2j]}, \quad 2 \leq j \leq n,$$

with

$$y_{[j,k]} := (s_j, s_{j+1}, \dots, s_k)^*, \quad 0 \leq j \leq k$$

and $y_{[j,k]} = 0_q$ if $j > k$. Define

$$u_{1,0} := 0_q, \quad u_{1,j} := \begin{pmatrix} 0_q \\ -y_{[0,j-1]} \end{pmatrix}, \quad u_{2,j} := -y_{[0,j]}, \quad 1 \leq j \leq n. \quad (16)$$

Let $\hat{H}_{1,j}$ (resp. $\hat{H}_{2,j}$) denote the Schur complement of the block Hankel matrices $H_{1,j-1}$ in $H_{1,j}$ (resp. $H_{2,j-1}$ in $H_{2,j}$) as seen in [45], [5, Equalities (2.5) and (2.6)]:

$$\begin{aligned} \hat{H}_{1,0} &:= s_0, & \hat{H}_{1,j} &:= s_{2j} - Y_{1,j}^* H_{1,j-1}^{-1} Y_{1,j}, \quad j \geq 1, \\ \hat{H}_{2,0} &:= s_1, & \hat{H}_{2,j} &:= s_{2j+1} - Y_{2,j}^* H_{2,j-1}^{-1} Y_{2,j}, \quad j \geq 1. \end{aligned}$$

These matrices are Hermitian positive definite matrices, and so are $H_{1,j}$ and $H_{2,j}$.

Now we partially reproduce left orthogonal matrix polynomials (OMP) on $[0, +\infty)$ [21].

Definition 3. Let σ be a $q \times q$ nonnegative measure on $[0, +\infty)$. A sequence $(P_j)_{j=0}^\infty$ of $q \times q$ matrix-valued polynomials is called orthogonal with respect to σ if

- i) each matrix-valued polynomial P_j is of degree j ;
- ii) the matrix-valued polynomials P_j satisfy the orthogonality relations:

$$\int_{[0, +\infty)} P_j(t) \sigma(dt) P_k^*(t) = \delta_{jk} C_{qj},$$

for all $j, k \in \mathbb{N}$. Here, δ_{jk} is the Kronecker delta, and C_{qj} is a constant nonnegative $q \times q$ matrix.

The following matrix polynomials seem to have been first introduced in [20].

Definition 4. Let $(s_j)_{j=0}^{2n}$ (resp. $(s_j)_{j=0}^{2n+1}$) be a Stieltjes positive sequence. Furthermore, let v_j , $u_{1,j}$, $u_{2,j}$ and R_j be as in (15), (16) and (14), respectively. Let

$$P_{1,0}(z) := I_q, \quad Q_{1,0}(z) := 0_q, \quad P_{2,0}(z) := I_q, \quad Q_{2,0}(z) := s_0.$$

For $j \geq 1$, set

$$\begin{aligned} P_{1,j}(z) &:= (-Y_{1,j}^* H_{1,j-1}^{-1}, I_q) R_j(z) v_j, \\ P_{2,j}(z) &:= (-Y_{2,j}^* H_{2,j-1}^{-1}, I_q) R_j(z) v_j, \\ Q_{1,j}(z) &:= -(-Y_{1,j}^* H_{1,j-1}^{-1}, I_q) R_j(z) u_{1,j}, \\ Q_{2,j}(z) &:= -(-Y_{2,j}^* H_{2,j-1}^{-1}, I_q) R_j(z) u_{2,j}. \end{aligned}$$

The matrix polynomials $Q_{1,j}$ and $Q_{2,j}$ are called *second-kind polynomials* with respect to $P_{1,j}$ and $P_{2,j}$, respectively.

In the next proposition, we recall that the matrix polynomials $P_{1,j}$ and $P_{2,j}$ are orthogonal with respect to positive measures supported on $[0, +\infty)$, and that the associated second-kind polynomials $Q_{1,j}$ and $Q_{2,j}$ can be expressed in terms of $P_{1,j}$ and $P_{2,j}$, respectively. For completeness, we also record a basic identity involving $P_{k,j}$ and $Q_{k,j}$.

Proposition 1. *a) The polynomials $P_{1,j}$ and $P_{2,j}$ are OMP with respect to σ and $t\sigma$, respectively. More precisely (as seen in [15, Remark 6.3]),*

$$\int_{[0,+\infty)} P_{k,j}(t)t^{k-1}\sigma(dt)P_{k,l}^*(t) = \begin{cases} 0_q & : j \neq l \\ \widehat{H}_{k,j} & : j = l \end{cases}, \quad k = 1, 2.$$

b) The following identities hold (as seen in [5, Remark D.6] and [5, Remark E.4]):

$$\begin{aligned} Q_{1,j}(x) &= \int_{[0,+\infty)} \frac{P_{1,j}(x) - P_{1,j}(t)}{x - t} \sigma(dt), \quad 0 \leq j \leq n, \\ Q_{2,j}(x) &= \int_{[0,+\infty)} \frac{xP_{2,j}(x) - tP_{2,j}(t)}{x - t} \sigma(dt), \quad 0 \leq j \leq n - 1. \end{aligned} \quad (17)$$

c) For $k = 1, 2$, the following identity holds (as seen in [5, Proposition 5.1 (a)])

$$P_{k,j}(z)Q_{k,j}^*(\bar{z}) - Q_{k,j}(z)P_{k,j}^*(\bar{z}) = 0_q.$$

The following proposition establishes a connection between HTM polynomials and orthogonal matrix polynomials of the first and second kinds. The representation below follows from [6, Theorem 6.1], while identities (19) and (20) are established in the proof of [6, Proposition 7.8].

Proposition 2. *For $n \geq 1$, every HTM polynomial admits the representation*

$$\mathbf{f}_n(z) = \begin{cases} (-1)^m(P_{1,m}^*(-\bar{z}^2) - zQ_{1,m}^*(-\bar{z}^2)), & n = 2m, \\ (-1)^m(Q_{2,m}^*(-\bar{z}^2) + zP_{2,m}^*(-\bar{z}^2)), & n = 2m + 1. \end{cases} \quad (18)$$

Moreover, the identities

$$\mathbf{h}_n(z) = (-1)^m P_{1,m}^*(-\bar{z}), \quad \mathbf{g}_n(z) = (-1)^{m+1} Q_{1,m}^*(-\bar{z}), \quad n = 2m, \quad (19)$$

and

$$\mathbf{h}_n(z) = (-1)^m Q_{2,m}^*(-\bar{z}), \quad \mathbf{g}_n(z) = (-1)^m P_{2,m}^*(-\bar{z}), \quad n = 2m + 1 \quad (20)$$

hold.

The matrix polynomials $P_{k,j}$ and $Q_{k,j}$ also appear in the representation of the extremal solutions of the TSMM problem.

In the proof of [6, Theorem 6.1], the so-called extremal solutions of the TSMM problem are represented by

$$G_{\sigma_{2m,\max}}(z) = -\frac{Q_{1,j}^*(\bar{z})}{P_{1,j}^*(\bar{z})} = \frac{\mathbf{g}_{2m}(-z)}{\mathbf{h}_{2m}(-z)} \quad (21)$$

and

$$G_{\sigma_{2m,\min}}(z) = -\frac{Q_{2,j}^*(\bar{z})}{zP_{2,j}^*(\bar{z})} = \frac{\mathbf{h}_{2m+1}(-z)}{(-z)\mathbf{g}_{2m+1}(-z)}. \quad (22)$$

Note that the matrix function $G_{\sigma_{2m,\max}}$ is defined for all $z \in \mathbb{C} \setminus [0, \infty)$ with $|z| > \eta_{2m}$, whereas $G_{\sigma_{2m,\min}}$ is defined for all $z \in \mathbb{C} \setminus [0, \infty)$ with $|z| > \rho_{2m+1}$. Here, η_{2m} and ρ_{2m+1} are defined in (8) and (9), respectively.

The Bezoutian and inertia for matrix polynomials

In this subsection, we reproduce some notions related to the inertia for a matrix polynomial [41, 42]. The inertia for scalar polynomials was considered in [40, page 462].

Let $L(\lambda) = L_0\lambda^\ell + L_1\lambda^{\ell-1} + \dots + \lambda L_{\ell-1} + L_\ell$ be a $q \times q$ matrix polynomial with coefficients $L_k \in \mathbb{C}^{q \times q}$. If $L_0 \neq 0_q$, then $\deg(L) = \ell$ is the degree of polynomial L . Furthermore, if $L_0 = I_q$, then L is called monic. Let M_1, L_1, M and L be polynomial matrices such that $M_1(\lambda)L_1(\lambda) - M(\lambda)L(\lambda) = 0$. The Bezoutian associated with the quadruple (M_1, L_1, M, L) is the block matrix

$$B_{M_1, M}(L, L_1) = (\Gamma_{j,k})_{j,k=1}^{m-1, \ell-1}. \quad (23)$$

The block entries $\Gamma_{j,k}$ of the quadruple (M_1, L_1, M, L) are determined by following equality:

$$\sum_{j,k=0}^{m-1, \ell-1} \Gamma_{j,k} \lambda^j \mu^k = \frac{1}{\lambda - \mu} (M_1(\lambda)L_1(\mu) - M(\lambda)L(\mu)). \quad (24)$$

The quantity ℓ (resp. m) denotes the maximal degree of $L(\lambda)$, $L_1(\lambda)$ (resp. $M(\lambda)$, $M_1(\lambda)$). Assume that there exists $\lambda_0 \in \mathbb{C}$ such that $\det L(\lambda_0) \neq 0$: on this case L is called regular.

Let L be a regular polynomial of degree ℓ . The triple

$$\gamma(L) = (\gamma_+(L), \gamma_-(L), \gamma_0(L)) \quad (25)$$

is called inertia with respect to the real axis, where $\gamma_+(L)$ (resp. $\gamma_-(L)$) is the number of zeros of $\det(L(\lambda))$ in the open upper (resp. open lower) half-plane, and $\gamma_0(L)$ is the number of zeros of $\det(L(\lambda))$ in the real axis. Now we reproduce [41, Corollary 2.2].

Corollary 1. *Let $L(\lambda)$ be a matrix polynomial with an invertible leading coefficient. The spectrum of $L(\lambda)$ lies in the upper half-plane if and only if there exists a regular matrix polynomial $L_1(\lambda)$ that satisfies*

$$L_1^*(\bar{\lambda})L_1(\lambda) = L^*(\bar{\lambda})L(\lambda)$$

such that the matrix

$$\frac{1}{i}B_{L_1^*, L^*}(L, L_1) \quad (26)$$

is positive definite.

Our strategy for analyzing the Hurwitzness of HTM polynomials $\mathbf{f}_n$ is based on the spectral symmetries of the quadruple

$$(\mathbf{f}_n^*(\bar{x}), \mathbf{f}_n(-x), \mathbf{f}_n^*(-\bar{x}), \mathbf{f}_n(x)).$$

The next lemma describes the spectral symmetries associated with this quadruple.

Lemma 1. *Let $F(z) = \sum_{k=0}^n A_k z^{n-k}$, $A_k \in \mathbb{C}^{q \times q}$, be a regular matrix polynomial, and let $F^*(z) := \sum_{k=0}^n A_k^* z^{n-k}$. If*

$$\sigma(F) := \{z \in \mathbb{C} : \det F(z) = 0\},$$

then $\sigma(F(-z)) = -\sigma(F)$, $\sigma(F^(\bar{z})) = \overline{\sigma(F)}$, and $\sigma(F^*(-\bar{z})) = -\overline{\sigma(F)}$.*

Proof. By definition, $\det F(-\lambda) = 0$ if and only if $-\lambda \in \sigma(F)$. Hence $\sigma(F(-z)) = -\sigma(F)$. Next, $\det F^*(\bar{z}) = \overline{\det F(z)}$. Therefore, $\det F^*(\bar{\lambda}) = 0$ if and only if $\det F(\lambda) = 0$. Consequently, $\sigma(F^*(\bar{z})) = \overline{\sigma(F)}$. Combining the previous two identities yields $\sigma(F^*(-\bar{z})) = -\overline{\sigma(F)}$.

Remark 2. *The four matrix polynomials $F(z)$, $F(-z)$, $F^*(\bar{z})$ and $F^*(-\bar{z})$ correspond to natural symmetries of the spectrum of F .*

Indeed, $F(-z)$ reflects the spectrum through the origin, $F^(\bar{z})$ reflects the spectrum across the real axis, and $F^*(-\bar{z})$ reflects the spectrum across the imaginary axis. Consequently, the Bezout form*

$$\frac{F^*(\bar{x})F(-y) - F^*(-\bar{x})F(y)}{x - y} \quad (27)$$

compares the spectrum of F with its reflection across the imaginary axis.

In the Hurwitz stable case, $\sigma(F) \subset \{z \in \mathbb{C} : \Re z < 0\}$, whereas $\sigma(F^(-\bar{z})) \subset \{z \in \mathbb{C} : \Re z > 0\}$. Hence these two spectra are separated by the imaginary axis. This separation is closely related to the definiteness properties of the associated Bezoutian.*

The generalized Anderson–Jury Bezoutian matrix

To compute the matrix (26) associated with an HTM polynomial, we use the generalized Bezoutian matrix and form introduced by Anderson and Jury in [2], which are defined for a $q \times q$ rational matrix function $W(z)$.

Suppose that $W(z)$ admits two matrix factorizations:

$$W(z) = A^{-1}(z)B(z) = D(z)C^{-1}(z).$$

The generalized Anderson–Jury Bezoutian form associated with the quadruple (A, B, C, D) is defined by

$$\Gamma(x, y) = \frac{1}{x - y} (A(x)D(y) - B(x)C(y)).$$

Using the notation (5), for HTM polynomials, the applicability of this construction follows from the fact that the $\mathbf{h}_n$ -part and the $\mathbf{g}_n$ -part of $\mathbf{f}_n$ determine the rational matrix functions

$$s_{\mathbf{f}_{2m}}(z) = \frac{\mathbf{g}_{2m}(-z)}{\mathbf{h}_{2m}(-z)} \quad (28)$$

and

$$s_{\mathbf{f}_{2m+1}}(z) = \frac{\mathbf{h}_{2m+1}(-z)}{(-z)\mathbf{g}_{2m+1}(-z)}. \quad (29)$$

Taking into account that the functions (28) and (29) (see also (21) and (22)) are Stieltjes transforms of positive matrix-valued measures; see (13), they satisfy the Hermitian symmetry relation

$$s_{\mathbf{f}_n}(z) = s_{\mathbf{f}_n}^*(\bar{z}). \quad (30)$$

Therefore, the Anderson–Jury Bezoutian construction can be applied. Consequently, rather than studying the Bezout form (27), we employ Bezoutian forms associated with the quadruples $(x\mathbf{g}_n^*, \mathbf{h}_n, \mathbf{h}_n^*, x\mathbf{g}_n)$ and $(\mathbf{g}_n^*, \mathbf{h}_n, \mathbf{h}_n^*, \mathbf{g}_n)$, corresponding to the cases $n = 2m + 1$ and $n = 2m$, respectively. See the definitions of $\mathcal{F}_n$, $\mathcal{G}_n^{(1)}$, and $\mathcal{G}_n^{(2)}$ in (31), (32), and (33), respectively, together with the relations between these forms given in (34).

3. Bezoutian and Hurwitz-type matrix polynomials

In this section, using the so-called matrix symmetrizers (50) and block Hankel matrices $H_{1,j}$ and $H_{2,j}$, we derive an explicit form of the Bezoutian for the quadruple $(\mathbf{f}_n^*(\bar{x}), \mathbf{f}_n(-x), \mathbf{f}_n^*(-\bar{x}), \mathbf{f}_n(x))$.

Instead of working directly with the Bezoutian form

$$\mathcal{F}_n(x, y) := \frac{\mathbf{f}_n^*(\bar{x})\mathbf{f}_n(-y) - \mathbf{f}_n^*(-\bar{x})\mathbf{f}_n(y)}{x - y}, \quad (31)$$

we use the Equality (2) and calculate the Bezoutian forms

$$\mathcal{G}_n^{(1)}(x, y) := \frac{1}{x^2 - y^2} (x^2 \mathbf{g}_n^*(\bar{x}^2) \mathbf{h}_n(y^2) - y^2 \mathbf{h}_n^*(\bar{x}^2) \mathbf{g}_n(y^2)) \quad (32)$$

and

$$\mathcal{G}_n^{(2)}(x, y) := \frac{xy}{x^2 - y^2} (\mathbf{g}_n^*(\bar{x}^2) \mathbf{h}_n(y^2) - \mathbf{h}_n^*(\bar{x}^2) \mathbf{g}_n(y^2)) \quad (33)$$

associated with the quadruples $(x\mathbf{g}_n^*, \mathbf{h}_n, \mathbf{h}_n^*, x\mathbf{g}_n)$ and $(\mathbf{g}_n^*, \mathbf{h}_n, \mathbf{h}_n^*, \mathbf{g}_n)$. Recall that a similar decomposition of the Bezoutian for scalar polynomials was previously studied; see [38, page 293] and [1, page 257]. Following [2] to compute the Bezoutian forms $\mathcal{G}_n^{(1)}(x, y)$ and $\mathcal{G}_n^{(2)}(x, y)$ we employ both representations (30) of the rational matrices (6) and (7).

It readily follows that

$$\mathcal{F}_n(x, y) = 2 \left(\mathcal{G}_n^{(1)}(x, y) + \mathcal{G}_n^{(2)}(x, y) \right). \quad (34)$$

In the next lemma, we demonstrate that the matrix functions $\mathcal{F}_n$, $\mathcal{G}_n^{(1)}$ and $\mathcal{G}_n^{(2)}$ are matrix polynomials in x and y . See [2, Remark 1].

Lemma 2. *Let $\mathcal{F}_n$, $\mathcal{G}_n^{(1)}$ and $\mathcal{G}_n^{(2)}$ be as defined in (31), (32) and (33), respectively. The functions $\mathcal{F}_n$, $\mathcal{G}_n^{(1)}$ and $\mathcal{G}_n^{(2)}$ are polynomials on x and y .*

Proof. Using the numerator on the right-hand side of (31), together with (18), and replacing y by x , we obtain

$$\begin{aligned} & \mathbf{f}_n(\bar{x})\mathbf{f}_n(-x) - \mathbf{f}_n(-\bar{x})\mathbf{f}_n(x) \\ &= 2x \begin{cases} P_{1,m}(-x^2)Q_{1,m}^*(-x^2) - Q_{1,m}(-x^2)P_{1,m}^*(-x^2), & n = 2m \\ P_{2,m}(-x^2)Q_{2,m}^*(-x^2) - Q_{2,m}(-x^2)P_{2,m}^*(-x^2), & n = 2m + 1 \end{cases} \\ &= 0. \end{aligned}$$

By [4, Chapter 1], any polynomial vanishing on the diagonal $x = y$ is divisible by $x - y$. Since $\mathbf{f}_n^*(\bar{x})\mathbf{f}_n(-y) - \mathbf{f}_n^*(-\bar{x})\mathbf{f}_n(y)$ has this property, $\mathcal{F}_n(x, y)$ is a polynomial. Arguing similarly with the numerators in (32) and (33), and following [4, Chapter 1], we show that $\mathcal{G}_n^{(1)}$ and $\mathcal{G}_n^{(2)}$ are polynomials in x and y .

For $k = 0, 1$, let

$$H_\infty^{(k)} := \begin{pmatrix} s_k & -s_{k+1} & s_{k+2} & \cdots \\ -s_{k+1} & s_{k+2} & \ddots & \\ & s_{k+2} & \ddots & \end{pmatrix} \quad (35)$$

be an infinite Hankel matrix.

A similar untruncated block Hankel matrix, denoted by $\mathbf{H}$, appears in the proof of [2, Lemma 2.3].

Lemma 3. *Let $\mathcal{F}_n$, $\mathcal{G}_n^{(1)}$ and $\mathcal{G}_n^{(2)}$ be as in (31), (32) and (33), respectively. Thus, for $n = 2m$, we have*

$$\mathcal{G}_{2m}^{(1)}(x, y) = \mathbf{h}_{2m}^*(\bar{x}^2) (x^{-2}I_q, x^{-4}I_q, \dots) H_\infty^{(1)} (\bar{y}^{-2}I_q, \bar{y}^{-4}I_q, \dots)^* \mathbf{h}_{2m}(y^2), \quad (36)$$

$$\mathcal{G}_{2m}^{(2)}(x, y) = -\mathbf{h}_{2m}^*(\bar{x}^2) (x^{-1}I_q, x^{-3}I_q, \dots) H_\infty^{(0)} (\bar{y}^{-1}I_q, \bar{y}^{-3}I_q, \dots)^* \mathbf{h}_{2m}(y^2). \quad (37)$$

For $n = 2m + 1$, we have

$$\mathcal{G}_{2m+1}^{(1)}(x, y) = \mathbf{g}_{2m+1}^*(\bar{x}^2) (I_q, x^{-2}I_q, \dots) H_{\infty}^{(0)} (I_q, \bar{y}^{-2}I_q, \dots)^* \mathbf{g}_{2m+1}(y^2), \quad (38)$$

$$\begin{aligned} \mathcal{G}_{2m+1}^{(2)}(x, y) = & -\mathbf{g}_{2m+1}^*(\bar{x}^2) (x^{-1}I_q, x^{-3}I_q, \dots) H_{\infty}^{(1)} (\bar{y}^{-1}I_q, \bar{y}^{-3}I_q, \dots)^* \\ & \cdot \mathbf{g}_{2m+1}(y^2). \end{aligned} \quad (39)$$

Proof. We prove (36). With (10), we have

$$\begin{aligned} \mathcal{G}_{2m}^{(1)}(x, y) &= \frac{1}{x^2 - y^2} \mathbf{h}_{2m}^*(\bar{x}^2) \left(x^2 \mathbf{h}_{2m}^{-1*}(\bar{x}^2) \mathbf{g}_{2m}^*(\bar{x}^2) - y^2 \mathbf{g}_{2m}(y^2) \mathbf{h}_{2m}^{-1}(y^2) \right) \\ &\quad \cdot \mathbf{h}_{2m}(y^2) \\ &= \frac{1}{x^2 - y^2} \mathbf{h}_{2m}^*(\bar{x}^2) (s_1(y^{-2} - x^{-2}) - s_2(y^{-4} - x^{-4}) + s_3(y^{-6} - x^{-6}) \\ &\quad - \dots) \mathbf{h}_{2m}(y^2) \\ &= \frac{1}{x^2 y^2} \mathbf{h}_{2m}^*(\bar{x}^2) (s_1 - s_2(x^{-2} + y^{-2}) + s_3(x^{-4} + x^{-2}y^{-2} + y^{-4}) \\ &\quad - \dots) \mathbf{h}_{2m}(y^2) \\ &= \mathbf{h}_{2m}^*(\bar{x}^2) (x^{-2}I_q, x^{-4}I_q, \dots) H_{\infty}^{(1)} (\bar{y}^{-2}I_q, \bar{y}^{-4}I_q, \dots)^* \mathbf{h}_{2m}(y^2). \end{aligned}$$

In the last equality, we substituted (35) for $k = 1$. Similarly, to prove (37), we use (10) and (35) for $k = 0$. Equalities (38) and (39) are proven by applying (11) and (35) for $k = 0$ and $k = 1$, respectively.

Remark 3. The infinite block Hankel matrix $H_{\infty}^{(1)}$ appearing in (36) arises naturally from the Laurent expansions in (10). Indeed, the expression

$$\frac{1}{x^2 - y^2} \left(x^2 \mathbf{h}_{2m}^{-1*}(\bar{x}^2) \mathbf{g}_{2m}^*(\bar{x}^2) - y^2 \mathbf{g}_{2m}(y^2) \mathbf{h}_{2m}^{-1}(y^2) \right)$$

generates an infinite series in negative powers of x and y , whose coefficients are precisely the moments $(s_j)_{j \geq 1}$, giving rise to the infinite block Hankel matrix $H_{\infty}^{(1)}$ in the representation

$$(x^{-2}I_q, x^{-4}I_q, \dots) H_{\infty}^{(1)} (\bar{y}^{-2}I_q, \bar{y}^{-4}I_q, \dots)^*.$$

However, since $\mathcal{G}_{2m}^{(1)}$ is a polynomial, only finitely many terms contribute to the products appearing in (36). Consequently, the right-hand side of (36) effectively reduces to a finite truncation of $H_{\infty}^{(1)}$.

The same observation applies to the right-hand sides of (37)–(39).

In view of Remark 3, the infinite block vector

$$(\bar{\xi}^{-2}I_q, \bar{\xi}^{-4}I_q, \dots)^* \mathbf{h}_{2m}(\xi^2)$$

can be decomposed into a polynomial part and a part containing strictly negative powers of ξ , namely,

$$(\bar{\xi}^{-2}I_q, \bar{\xi}^{-4}I_q, \dots)^* \mathbf{h}_{2m}(\xi^2) = \mathbf{h}_{2m}^{(11)}(\xi) + \mathbf{h}_{2m}^{(12)}(\xi). \quad (40)$$

Similarly, we have

$$(\bar{\xi}^{-1}I_q, \bar{\xi}^{-3}I_q, \dots)^* \mathbf{h}_{2m}(\xi^2) = \mathbf{h}_{2m}^{(21)}(\xi) + \mathbf{h}_{2m}^{(22)}(\xi), \quad (41)$$

$$(I_q, \bar{\xi}^{-2}I_q, \dots)^* \mathbf{g}_{2m+1}(\xi^2) = \mathbf{g}_{2m+1}^{(11)}(\xi) + \mathbf{g}_{2m+1}^{(12)}(\xi), \quad (42)$$

$$(\bar{\xi}^{-1}I_q, \bar{\xi}^{-3}I_q, \dots)^* \mathbf{g}_{2m+1}(\xi^2) = \mathbf{g}_{2m+1}^{(21)}(\xi) + \mathbf{g}_{2m+1}^{(22)}(\xi). \quad (43)$$

Here, $\mathbf{h}_{2m}^{(k1)}(\xi)$ (resp. $\mathbf{g}_{2m+1}^{(k1)}(\xi)$), for $k = 1, 2$, are polynomials in ξ , whereas $\mathbf{h}_{2m}^{(k2)}(\xi)$ (resp. $\mathbf{g}_{2m+1}^{(k2)}(\xi)$), for $k = 1, 2$, are polynomials in ξ^{-1} .

For a natural number m , let

$$\tilde{J}_m := \text{diag} (I_q, (-1)I_q, (-1)^2I_q, \dots, (-1)^m I_q) \quad (44)$$

be a $(m+1)q \times (m+1)q$ matrix.

The following remark explains how the infinite Hankel matrices $H_\infty^{(k)}$ reduce to finite block Hankel matrices when acting on the polynomial components $\mathbf{h}_{2m}^{(k1)}(\xi)$ (resp. $\mathbf{g}_{2m+1}^{(k1)}(\xi)$), for $k = 1, 2$, introduced in (40)–(43). This follows from direct calculations based on the corresponding decompositions.

Remark 4. Let $\mathbf{h}_{2m}^{(11)}(\xi)$, $\mathbf{h}_{2m}^{(21)}(\xi)$, $\mathbf{g}_{2m+1}^{(11)}(\xi)$ and $\mathbf{g}_{2m+1}^{(21)}(\xi)$ be as defined in (40), (41), (42) and (43), respectively. Furthermore, let $H_{1,j}$, $H_{2,j}$ and $J_m^{(k)}$ be as defined in (12) and (44), respectively. Thus, the following equalities hold

$$\mathbf{h}_{2m}^{(11)*}(\bar{x})H_\infty^{(1)}\mathbf{h}_{2m}^{(11)}(y) = \widehat{\mathbf{h}}_{2m}^{(11)*}(\bar{x})\tilde{J}_{m-1}H_{2,m-1}\tilde{J}_{m-1}\widehat{\mathbf{h}}_{2m}^{(11)}(y), \quad (45)$$

$$\mathbf{h}_{2m}^{(21)*}(\bar{x})H_\infty^{(1)}\mathbf{h}_{2m}^{(21)}(y) = \widehat{\mathbf{h}}_{2m}^{(21)*}(\bar{x})\tilde{J}_{m-1}H_{1,m-1}\tilde{J}_{m-1}\widehat{\mathbf{h}}_{2m}^{(21)}(y), \quad (46)$$

$$\mathbf{g}_{2m+1}^{(11)*}(\bar{x})H_\infty^{(1)}\mathbf{g}_{2m+1}^{(11)}(y) = \widehat{\mathbf{g}}_{2m+1}^{(11)*}(\bar{x})\tilde{J}_mH_{1,m}\tilde{J}_m\widehat{\mathbf{g}}_{2m+1}^{(11)}(y), \quad (47)$$

$$\mathbf{g}_{2m+1}^{(21)*}(\bar{x})H_\infty^{(1)}\mathbf{g}_{2m+1}^{(21)}(y) = \widehat{\mathbf{g}}_{2m+1}^{(21)*}(\bar{x})\tilde{J}_{m-1}H_{2,m-1}\tilde{J}_{m-1}\widehat{\mathbf{g}}_{2m+1}^{(21)}(y), \quad (48)$$

where

$$\widehat{\mathbf{h}}_{2m}^{(11)}, \widehat{\mathbf{h}}_{2m}^{(21)}, \widehat{\mathbf{g}}_{2m+1}^{(11)} \quad \text{and} \quad \widehat{\mathbf{g}}_{2m+1}^{(21)}$$

represent finite-dimensional restrictions of the matrices $\mathbf{h}_{2m}^{(11)}$, $\mathbf{h}_{2m}^{(21)}$, $\mathbf{g}_{2m+1}^{(11)}$ and $\mathbf{g}_{2m+1}^{(21)}$. In particular, the polynomial part of

$$\widehat{\mathbf{h}}_{2m}^{(11)}(y) = (\bar{y}^{-2}\mathbf{h}_{2m}^*(\bar{y}^2), \bar{y}^{-4}\mathbf{h}_{2m}^*(\bar{y}^2), \dots, \bar{y}^{-2m}\mathbf{h}_{2m}^*(\bar{y}^2))^* \quad (49)$$

is the restriction of the polynomial part of the matrix

$$\mathbf{h}_{2m}^{(11)}(y) = (\bar{y}^{-2}\mathbf{h}_{2m}^*(\bar{y}^2), \bar{y}^{-4}\mathbf{h}_{2m}^*(\bar{y}^2), \dots, \bar{y}^{-2m}\mathbf{h}_{2m}^*(\bar{y}^2), 0_q, 0_q, \dots)^*.$$

Equalities (45)–(48) correspond to those in the proof of [2, Lemma 2.3], where, instead of the matrices $H_{1,j}$ and $H_{2,j}$, the truncated block Hankel matrix $\mathbf{H}_{nm}$ is used. Moreover, in place of the matrix $H_{\infty}^{(1)}$, the infinite Hankel matrix H is employed.

For a matrix polynomial $Q(x) = Q_0x^\ell + Q_1x^{\ell-1} + \dots + Q_{\ell-1}x + Q_\ell$, the Hankel block matrix

$$S(Q) := \begin{pmatrix} Q_{\ell-1} & Q_{\ell-2} & \dots & Q_1 & Q_0 \\ Q_{\ell-2} & Q_{\ell-3} & \dots & Q_0 & 0_q \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ Q_0 & 0_q & \dots & 0_q & 0_q \end{pmatrix} \quad (50)$$

is called a matrix symmetrizer of $Q(x)$. This matrix provides a finite-dimensional representation of the coefficients of Q and will be used to express the Bezoutian in terms of block Hankel matrices. In [40, page 455], the concept of symmetrizer is introduced for scalar polynomials.

Additionally, let

$$F_0^{2k}(x) := \left(I_q, \bar{x}^2 I_q, \dots, \bar{x}^{2k} I_q \right)^*, \quad (51)$$

$$F_1^{2k-1}(x) := \left(\bar{x} I_q, \bar{x}^3 I_q, \dots, \bar{x}^{2k-1} I_q \right)^*. \quad (52)$$

In the following theorem, added by symmetrizers of corresponding matrices and the block Hankel matrices $H_{k,j}$, we obtain a factorized expression of the Bezoutian forms $\mathcal{G}_{2m}^{(1)}$, $\mathcal{G}_{2m}^{(2)}$, $\mathcal{G}_{2m+1}^{(1)}$ and $\mathcal{G}_{2m+1}^{(2)}$.

Theorem 1. Let $\mathcal{G}_{2m}^{(k)}$, $\mathcal{G}_{2m+1}^{(k)}$ for $k = 1, 2$ be as defined in (32) and (33). Additionally, let $\mathbf{h}_n$, $\mathbf{g}_n$, S , $\tilde{J}_m$, $H_{1,j}$, $H_{2,j}$, $F_0^{2k}(x)$ and $F_1^{2k-1}(x)$ be as in (3), (4), (50), (44), (12), (51) and (52), respectively.

Therefore, for $n = 2m$, the following equality holds:

$$\mathcal{G}_{2m}^{(1)}(x, y) = F_0^{(2m-2)*}(\bar{x}) S(\mathbf{h}_{2m}^*) \tilde{J}_{m-1} H_{2,m-1} \tilde{J}_{m-1} S(\mathbf{h}_{2m}) F_0^{(2m-2)}(y), \quad (53)$$

$$\mathcal{G}_{2m}^{(2)}(x, y) = -F_1^{(2m-1)*}(\bar{x}) S(\mathbf{h}_{2m}^*) \tilde{J}_{m-1} H_{1,m-1} \tilde{J}_{m-1} S(\mathbf{h}_{2m}) F_1^{(2m-1)}(y). \quad (54)$$

For $n = 2m + 1$, we have

$$\mathcal{G}_{2m+1}^{(1)}(x, y) = F_0^{(2m)*}(\bar{x}) S(\mathbf{g}_{2m+1}^*) \tilde{J}_m H_{1,m} \tilde{J}_m S(\mathbf{g}_{2m+1}) F_0^{(2m)}(y). \quad (55)$$

Moreover,

$$\mathcal{G}_{2m+1}^{(2)}(x, y) = -F_1^{(2m-1)*}(\bar{x}) S(\tilde{\mathbf{g}}_{2m+1}^*) \tilde{J}_{m-1} H_{2,m-1} \tilde{J}_{m-1} S(\tilde{\mathbf{g}}_{2m+1}) F_1^{(2m-1)}(y), \quad (56)$$

where

$$\tilde{\mathbf{g}}_{2m+1}(z) := I_q z^{m-1} + A_2 z^{m-2} + \dots + A_{2m-2}. \quad (57)$$

Proof. We prove (53). From (36), (40) and (45), we have

$$\mathcal{G}_{2m}^{(1)}(x, y) = \widehat{\mathbf{h}}_{2m}^{(11)*}(\bar{x}) \widetilde{J}_{m-1} H_{2,m-1} \widetilde{J}_{m-1} \widehat{\mathbf{h}}_{2m}^{(11)}(y). \quad (58)$$

We examine $\widehat{\mathbf{h}}_{2m}^{(11)}(y)$. By using (49) and (50), we have

$$\begin{aligned} \widehat{\mathbf{h}}_{2m}^{(11)}(y) &= \begin{pmatrix} A_{2m-2} + \dots + A_2 y^{2m-4} + I_q y^{2m-2} \\ A_{2m-4} + \dots + A_2 y^{2m-6} + I_q y^{2m-4} \\ \vdots \\ I_q \end{pmatrix} \\ &= \begin{pmatrix} A_{2m-2} & \dots & A_2 & I_q \\ A_{2m-4} & \dots & I_q & \\ \vdots & & & \\ I_q & & & \end{pmatrix} \begin{pmatrix} I_q \\ y^2 I_q \\ \vdots \\ y^{2m-2} I_q \end{pmatrix} \\ &= S(\mathbf{h}_{2m}) (I_q, \bar{y}^2 I_q, \dots, \bar{y}^{2m-2} I_q)^*. \end{aligned} \quad (59)$$

By substituting (59) for (58), we obtain (53). Similarly, by using

$$\mathcal{G}_{2m}^{(2)}(x, y) = \widehat{\mathbf{h}}_{2m}^{(21)*}(\bar{x}) \widetilde{J}_{m-1} H_{1,m-1} \widetilde{J}_{m-1} \widehat{\mathbf{h}}_{2m}^{(21)}(y), \quad (60)$$

$$\mathcal{G}_{2m+1}^{(1)}(x, y) = \widehat{\mathbf{g}}_{2m+1}^{(11)*}(\bar{x}) \widetilde{J}_m H_{1,m-1} \widetilde{J}_m \widehat{\mathbf{g}}_{2m+1}^{(11)}(y), \quad (61)$$

$$\mathcal{G}_{2m+1}^{(2)}(x, y) = \widehat{\mathbf{g}}_{2m+1}^{(21)*}(\bar{x}) \widetilde{J}_{m-1} H_{2,m-1} \widetilde{J}_{m-1} \widehat{\mathbf{g}}_{2m+1}^{(21)}(y), \quad (62)$$

one can prove (54), (55) and (56). Equalities (60), (61) and (62) readily follow from (46), (47), (48) and (41), as well as from (42) and (43). Additionally, one also uses (37), (38), (39), (51) and (52).

Remark 5. The polynomial $\widetilde{\mathbf{g}}_{2m+1}$ defined in (57) has degree $m-1$, in contrast to the original polynomial $\mathbf{g}_{2m+1}$ defined in (4). Using $\widetilde{\mathbf{g}}_{2m+1}$, we construct the symmetrizer $S(\widetilde{\mathbf{g}}_{2m+1})$ appearing in the matrix $\mathcal{W}_{2m+1}^{(0)}$ in (76). This is consistent with the fact that the block Hankel matrices $H_{1,m}$ and $H_{2,m-1}$ have different dimensions.

In the following corollary, we represent $\mathcal{F}_n$ as a matrix quadratic form regarding the block Hankel matrices $H_{1,j}$ and $H_{2,j}$, as well as the symmetrizer matrices $S(\mathbf{h}_{2m})$, $S(\mathbf{g}_{2m+1})$ and $S(\widetilde{\mathbf{g}}_{2m+1})$.

Corollary 2. Assume the same conditions and notations as in Theorem 1. Let $\widetilde{J}_m$ be as in (44). Moreover, let

$$\mathcal{J}_r^{(\ell)} := \begin{pmatrix} \widetilde{J}_\ell & 0 \\ 0 & \widetilde{J}_r \end{pmatrix} \quad (63)$$

for all $r, \ell > 1$. Furthermore, for $k = 0, 1$ let

$$\mathcal{W}_{2m}^{(k)} := \begin{pmatrix} S(\mathbf{h}_{2m}^*) & 0 \\ 0 & S(\mathbf{h}_{2m}^*) \end{pmatrix} \mathcal{J}_{m-1}^{(m-1)} \begin{pmatrix} H_{2,m-1} & 0 \\ 0 & (-1)^k H_{1,m-1} \end{pmatrix} \mathcal{J}_{m-1}^{(m-1)} \cdot \begin{pmatrix} S(\mathbf{h}_{2m}) & 0 \\ 0 & S(\mathbf{h}_{2m}) \end{pmatrix}, \quad (64)$$

$$\mathcal{W}_{2m+1}^{(k)} := \begin{pmatrix} S(\mathbf{g}_{2m+1}^*) & 0 \\ 0 & S(\mathbf{g}_{2m+1}^*) \end{pmatrix} \mathcal{J}_{m-1}^{(m)} \begin{pmatrix} H_{1,m} & 0 \\ 0 & (-1)^k H_{2,m-1} \end{pmatrix} \mathcal{J}_{m-1}^{(m)} \cdot \begin{pmatrix} S(\mathbf{g}_{2m+1}) & 0 \\ 0 & S(\mathbf{g}_{2m+1}) \end{pmatrix}. \quad (65)$$

With $\mathcal{F}_n$ be as in (31), the following equalities hold:

$$\mathcal{F}_{2m}(x, y) = 2 \begin{pmatrix} F_0^{(2m-2)}(\bar{x}) \\ F_1^{(2m-1)}(\bar{x}) \end{pmatrix}^* \mathcal{W}_{2m}^{(1)} \begin{pmatrix} F_0^{(2m-2)}(y) \\ F_1^{(2m-1)}(y) \end{pmatrix}, \quad (66)$$

and

$$\mathcal{F}_{2m+1}(x, y) = 2 \begin{pmatrix} F_0^{(2m)}(\bar{x}) \\ F_1^{(2m-1)}(\bar{x}) \end{pmatrix}^* \mathcal{W}_{2m+1}^{(1)} \begin{pmatrix} F_0^{(2m)}(y) \\ F_1^{(2m-1)}(y) \end{pmatrix}. \quad (67)$$

Proof. Equality (66) follows directly from (34), (53), (54) and (64). In a similar manner, Equality (67) can readily be derived from (34), (55), (56) and (65).

4. Hurwitzness of Hurwitz-type matrix polynomials

In this section, using the explicit form of the Bezoutian matrix (66) and (67), we verify that an HTM polynomial $\mathbf{f}_n$ is indeed a Hurwitz matrix polynomial.

In [49], the Hurwitz property of HTM polynomials was studied. However, the proof of [49, Theorem 3.5] concerning the inertia of the matrix polynomial $\mathbf{f}_n$ for $n = 2m$ is neither explicit nor complete, and a proof for the case $n = 2m + 1$ is not provided. See Remark 14.

Recall that the inertia of matrix polynomials, denoted by $\gamma(\mathbf{f}_n)$, provides the number of eigenvalues of $\mathbf{f}_n$ with positive, negative and real zero parts, respectively; see [41, page 410].

We rewrite Corollary 2 for the pair (ix, iy) instead of (x, y) .

Corollary 3. Assume the same conditions and notations as in Theorem 1. Moreover, let $\mathcal{F}_n$, $\mathcal{J}_k^\ell$, $\mathcal{W}_{2m}^{(0)}$ and $\mathcal{W}_{2m+1}^{(0)}$ be as in (31), (63), (64) and (65), respectively. Thus, the following equalities hold:

$$\mathcal{F}_{2m}(ix, iy) = 2 \begin{pmatrix} F_0^{(2m-2)}(\bar{x}) \\ F_1^{(2m-1)}(\bar{x}) \end{pmatrix}^* \mathcal{J}_{m-1}^{(m-1)} \mathcal{W}_{2m}^{(0)} \mathcal{J}_{m-1}^{(m-1)} \begin{pmatrix} F_0^{(2m-2)}(y) \\ F_1^{(2m-1)}(y) \end{pmatrix} \quad (68)$$

and

$$\mathcal{F}_{2m+1}(ix, iy) = 2 \begin{pmatrix} F_0^{(2m)}(\bar{x}) \\ F_1^{(2m-1)}(\bar{x}) \end{pmatrix}^* \mathcal{J}_{m-1}^{(m)} \mathcal{W}_{2m+1}^{(0)} \mathcal{J}_{m-1}^{(m)} \begin{pmatrix} F_0^{(2m)}(y) \\ F_1^{(2m-1)}(y) \end{pmatrix}. \quad (69)$$

Proof. Equalities (68) and (69) readily follow from (66) and (67), respectively. Let

$$\begin{aligned} V_{[0,2m-1]}(x) &:= (I_q, \bar{x}I_q, \bar{x}^2I_q, \dots, \bar{x}^{2m-2}I_q, \bar{x}^{2m-1}I_q)^*, \\ V_{[0,2m]}(x) &:= (I_q, \bar{x}I_q, \bar{x}^2I_q, \dots, \bar{x}^{2m-1}I_q, \bar{x}^{2m}I_q)^*. \end{aligned}$$

Furthermore, let $\mathcal{T}_{2m}$ (resp. $\mathcal{T}_{2m+1}$) be elementary block matrices such that

$$\begin{pmatrix} F_0^{(2m-2)}(x) \\ F_1^{(2m-1)}(x) \end{pmatrix} = \mathcal{J}_{m-1}^{(m-1)} \mathcal{T}_{2m} V_{[0,2m-1]}(x) \quad (70)$$

and

$$\begin{pmatrix} F_0^{(2m)}(x) \\ F_1^{(2m-1)}(x) \end{pmatrix} = \mathcal{J}_{m-1}^{(m)} \mathcal{T}_{2m+1} V_{[0,2m]}(x). \quad (71)$$

The following remark readily follows.

Remark 6. Using (70) and (71), equalities (68) and (69) can be written as follows:

$$\mathcal{F}_{2m}(ix, iy) = 2V_{[0,2m-1]}^*(\bar{x}) \mathcal{T}_{2m}^* \mathcal{J}_{m-1}^{(m-1)} \mathcal{W}_{2m}^{(0)} \mathcal{J}_{m-1}^{(m-1)} \mathcal{T}_{2m} V_{[0,2m-1]}(y), \quad (72)$$

$$\mathcal{F}_{2m+1}(ix, iy) = 2V_{[0,2m]}^*(\bar{x}) \mathcal{T}_{2m+1}^* \mathcal{J}_{m-1}^{(m)} \mathcal{W}_{2m+1}^{(0)} \mathcal{J}_{m-1}^{(m)} \mathcal{T}_{2m+1} V_{[0,2m]}(y). \quad (73)$$

The positivity of the matrices $\mathcal{W}_{2m}^{(0)}$ and $\mathcal{W}_{2m+1}^{(0)}$ is consistent with the spectral separation described in Remark 2. Indeed, the substitution $z = i\lambda$ transforms separation with respect to the imaginary axis into a Hermitian positivity structure for the associated Bezoutian kernel.

We now state the main result of the present work. The key step is to express the matrix

$$\frac{1}{i} B_{L_1^*, L^*}(L, L_1)$$

in terms of the Bezoutian forms obtained above and to prove its positive definiteness. This allows us to establish the Hurwitz property of HTM polynomials.

Theorem 2. *Each HTM polynomial is a Hurwitz matrix polynomial.*

Proof We use Corollary 1 with

$$L(\lambda) = \mathbf{f}_n(i\lambda) \quad \text{and} \quad L_1(\lambda) = \mathbf{f}_n(-i\lambda) \quad (74)$$

and Equalities (23), (24), (31), (32), (33) and (34). For $n = 2m$, by using (72), we have

$$\frac{1}{i} B_{L_1^*, L^*}(L, L_1) = 2\mathcal{T}_{2m}^* \mathcal{J}_{m-1}^{(m-1)} \mathcal{W}_{2m}^{(0)} \mathcal{J}_{m-1}^{(m-1)} \mathcal{T}_{2m}. \quad (75)$$

By employing (73) for $n = 2m + 1$, we have

$$\frac{1}{i} B_{L_1^*, L^*}(L, L_1) = 2\mathcal{T}_{2m+1}^* \mathcal{J}_{m-1}^{(m)} \mathcal{W}_{2m+1}^{(0)} \mathcal{J}_{m-1}^{(m)} \mathcal{T}_{2m+1}. \quad (76)$$

The right-hand sides of (75) and (76) are positive definite matrices because

$$\begin{pmatrix} H_{2,m-1} & 0 \\ 0 & H_{1,m-1} \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} H_{1,m} & 0 \\ 0 & H_{2,m-1} \end{pmatrix} \quad (77)$$

are positive definite, and both $\mathcal{J}_{m-1}^{(m-1)} \mathcal{T}_{2m}$ and $\mathcal{J}_{m-1}^{(m)} \mathcal{T}_{2m+1}$ are invertible matrices.

By Corollary 1, the spectra of

$$L(\lambda) = \mathbf{f}_n(i\lambda)$$

belong to the upper half-plane of $\mathbb{C}$. Consequently, the spectra of $\mathbf{f}_n(z)$ lie in the open left half-plane of $\mathbb{C}$. Hence, the HTM polynomials $\mathbf{f}_n$ are Hurwitz matrix polynomials.

Remark 7. By employing (25), note that

$$\gamma_-(\mathbf{f}_n(i\lambda)) = \gamma_0(\mathbf{f}_n(i\lambda)) = 0,$$

and therefore

$$\gamma(\mathbf{f}_n(i\lambda)) = \gamma_+(\mathbf{f}_n(i\lambda)).$$

This is equivalent to the fact that all zeros of $\det \mathbf{f}_n(\lambda)$ lie in the open left half-plane.

The following remark illustrates the previous observation in the simplest nontrivial case.

Remark 8. A $q \times q$ HTM polynomial $\mathbf{f}_2$ of degree 2 is a Hurwitz matrix polynomial; it admits the representation

$$\mathbf{f}_2(z) = I_q z^2 + s_0 z + s_0^{-1} s_1.$$

The next remark concerns the matrix measures associated with HTM polynomials and their connection with the corresponding orthogonal matrix polynomials.

Remark 9. A $q \times q$ matrix $\tilde{\sigma}(\cdot)$ defined on $[0, \infty)$ is called monotonically nondecreasing if $\tilde{\sigma}(t)$ is self-adjoint for all $t \in [0, \infty)$ and $\tilde{\sigma}(t_2) - \tilde{\sigma}(t_1) \geq 0$ for $t_1 \leq t_2$.

For $n = 2m + 1$ (resp. $n = 2m$), the points of discontinuity of the distribution function $\tilde{\sigma}_{2m,\max}$ (resp. $\tilde{\sigma}_{2m,\min}$) that correspond to the measure $\sigma_{2m,\max}$ (resp. $\sigma_{2m,\min}$) are given by the roots of the polynomial $\det(P_{1,m}^*(-\bar{z}))$ (resp. $\det(zP_{2,m}^*(-\bar{z}))$); see (21) and (22).

On the other hand, the polynomials (19) and (20), which are constructed using the moments s_j , can be obtained from an absolutely continuous distribution on $[0, \infty)$ that corresponds to a positive measure on $[0, \infty)$, as shown in the next example.

The following example illustrates the previous construction in the case of a fifth-degree HTM polynomial. For the matrix polynomial $\mathbf{f}_5$, we compute the Bezoutian forms $\mathcal{G}_5^{(1)}$ and $\mathcal{G}_5^{(2)}$, the symmetrizers $S(\mathbf{g}_5)$ and $S(\tilde{\mathbf{g}}_5)$, the matrices $\tilde{\mathcal{J}}_1, \tilde{\mathcal{J}}_2, \mathcal{J}_1^{(2)}$, as well as the matrices $\mathcal{W}_5^{(0)}$ and $\mathcal{T}_5$. Using these matrices, we obtain an explicit representation of

$$\frac{1}{i}B_{L_1^*, L^*}(L, L_1),$$

whose positive definiteness ensures that $\mathbf{f}_5$ is a Hurwitz matrix polynomial.

Example 1. *Let*

$$\mathbf{f}_5(z) = z^5 I_2 + A_1 z^4 + A_2 z^3 + A_3 z^2 + A_4 z + A_5, \quad (78)$$

$$\text{where } A_1 = \begin{pmatrix} 2 & -i \\ i & 1 \end{pmatrix}, A_2 = \begin{pmatrix} \frac{486}{37} & 0 \\ -\frac{264}{37}i & 6 \end{pmatrix}, A_3 = \begin{pmatrix} \frac{560}{37} & -5i \\ 5i & 5 \end{pmatrix},$$

$$A_4 = \begin{pmatrix} \frac{1158}{37} & 0 \\ -\frac{936}{37}i & 6 \end{pmatrix} \text{ and } A_5 = \begin{pmatrix} \frac{292}{37} & -2i \\ 2i & 2 \end{pmatrix}.$$

The matrix polynomials $\mathbf{h}_5$ and $\mathbf{g}_5$ have the form

$$\mathbf{h}_5(x) = A_1 x^2 + A_3 x + A_5, \quad \text{and} \quad \mathbf{g}_5(x) = I_2 x^2 + A_2 x + A_4 \quad (79)$$

Using (79) together with the Laurent expansions (10) and (11), we obtain the first four moments:

$$s_0 = \begin{pmatrix} 2 & -i \\ i & 1 \end{pmatrix}, \quad s_1 = \begin{pmatrix} 4 & -i \\ i & 1 \end{pmatrix}, \quad s_2 = \begin{pmatrix} 16 & -2i \\ 2i & 2 \end{pmatrix}, \quad (80)$$

$$s_3 = \begin{pmatrix} 96 & -6i \\ 6i & 6 \end{pmatrix}, \quad s_4 = \begin{pmatrix} 768 & -24i \\ 24i & 24 \end{pmatrix}. \quad (81)$$

The Bezoutian forms $\mathcal{G}_5^{(1)}$ and $\mathcal{G}_5^{(2)}$ defined in (32) and (33) are given by

$$\mathcal{G}_5^{(1)}(x, y) = \begin{pmatrix} g_{11}(x, y) & -i g_{12}(x, y) \\ i g_{12}(x, y) & g_{22}(x, y) \end{pmatrix}$$

where

$$g_{11} = \frac{2}{1369} (37x^4 (37y^4 + 280y^2 + 146) + 4x^2 (2590y^4 + 22883y^2 + 15297) + 5402y^4 + 61188y^2 + 134436)$$

$$g_{12} = ((y^4 + 5y^2 + 2)x^4 + (5y^4 + 26y^2 + 12)x^2 + 2(y^4 + 6y^2 + 6))$$

$$g_{22} = (y^4 + 5y^2 + 2)x^4 + (5y^4 + 26y^2 + 12)x^2 + 2(y^4 + 6y^2 + 6)$$

and

$$\mathcal{G}_5^{(2)}(x, y) = \begin{pmatrix} -\frac{4xy(37(37y^2+272)x^2+10064y^2+88236)}{1369} & ixy((y^2+4)x^2+4y^2+18) \\ -ixy((y^2+4)x^2+4y^2+18) & -xy((y^2+4)x^2+4y^2+18) \end{pmatrix}$$

The matrices $S(\mathbf{g}_5)$ and $S(\tilde{\mathbf{g}}_5)$, defined in (50), have the form

$$S(\mathbf{g}_5) = \begin{pmatrix} A_4 & A_2 & I_2 \\ A_2 & I_2 & 0_2 \\ I_2 & 0_2 & 0_2 \end{pmatrix} = \begin{pmatrix} \frac{1158}{37} & 0 & \frac{486}{37} & 0 & 1 & 0 \\ -\frac{936i}{37} & 6 & -\frac{264i}{37} & 6 & 0 & 1 \\ \frac{486}{37} & 0 & 1 & 0 & 0 & 0 \\ -\frac{264i}{37} & 6 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix} \quad (82)$$

and

$$S(\tilde{\mathbf{g}}_5) = \begin{pmatrix} A_2 & I_2 \\ I_2 & 0_2 \end{pmatrix} = \begin{pmatrix} \frac{1158}{37} & 0 & \frac{486}{37} & 0 & 1 & 0 \\ -\frac{936i}{37} & 6 & -\frac{264i}{37} & 6 & 0 & 1 \\ \frac{486}{37} & 0 & 1 & 0 & 0 & 0 \\ -\frac{264i}{37} & 6 & 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \end{pmatrix}. \quad (83)$$

The matrix polynomial $\tilde{\mathbf{g}}_5$ has the form

$$\tilde{\mathbf{g}}_5(x) = I_2 x + A_2.$$

Furthermore, the matrices $F_0^{(4)}$ and $F_1^{(3)}$, defined in (51) and (52), have the form

$$F_0^{(4)}(x) = (I_2 \quad \bar{x}^2 I_2 \quad \bar{x}^4 I_2)^* \quad \text{and} \quad F_1^{(3)}(x) = (I_2 \quad \bar{x}^2 I_2)^*. \quad (84)$$

The matrices $\tilde{J}_2$ and $\tilde{J}_1$, introduced in (44), take the form

$$\tilde{J}_2 = \begin{pmatrix} I_2 & 0_2 & 0_2 \\ 0_2 & -I_2 & 0_2 \\ 0_2 & 0_2 & 1_2 \end{pmatrix}, \quad \text{and} \quad \tilde{J}_1 = \begin{pmatrix} I_2 & 0_2 \\ 0_2 & -I_2 \end{pmatrix}. \quad (85)$$

Using (80) and (81), we construct the matrices $H_{1,2}$ and $H_{2,1}$. One readily verifies that these block matrices are positive definite. From (82), (83), (84), and (85), we obtain the right-hand sides of (55) and (56), which allow us to conclude that $\mathcal{G}_5^{(1)}$ and $-\mathcal{G}_5^{(2)}$ are positive quadratic forms.

The matrix $\frac{1}{i} B_{L_1^*, L^*}(L, L_1)$ in (76) can be written as the product of the matrices

$$\mathcal{J}_1^{(2)} = \begin{pmatrix} \tilde{J}_2 & 0 \\ 0 & \tilde{J}_1 \end{pmatrix},$$

$$\mathcal{W}_5^{(0)} = \begin{pmatrix} S(\mathbf{g}_5^*) & 0 \\ 0 & S(\tilde{\mathbf{g}}_5^*) \end{pmatrix} \mathcal{J}_1^{(2)} \begin{pmatrix} H_{1,2} & 0 \\ 0 & H_{2,1} \end{pmatrix} \mathcal{J}_1^{(2)} \begin{pmatrix} S(\mathbf{g}_5) & 0 \\ 0 & S(\tilde{\mathbf{g}}_5) \end{pmatrix}$$

and the matrix $\mathcal{T}_5$, which appears in (73), has the form

$$\mathcal{T}_5 = \begin{pmatrix} I_2 & 0_2 & 0_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & 0_2 & I_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & 0_2 & 0_2 & I_2 \\ 0_2 & I_2 & 0_2 & 0_2 & 0_2 & 0_2 \\ 0_2 & 0_2 & 0_2 & 0_2 & I_2 & 0_2 \end{pmatrix}.$$

From these matrices, together with (73), (76), Corollary 1, and the second matrix in (77), it follows that the matrix polynomial $\mathbf{f}_5$ is a Hurwitz matrix polynomial.

The following remark illustrates how HTM polynomials can be generated from matrix-valued measures on $[0, \infty)$.

Remark 10. *The polynomial $\mathbf{f}_5$ in (78) can be constructed starting from the matrix-valued distribution*

$$\tilde{\sigma}(x) = \begin{pmatrix} 4 - 2e^{-x/2} & ie^{-x} \\ -ie^{-x} & 2 - e^{-x} \end{pmatrix},$$

defined on $[0, \infty)$.

More generally, the construction of HTM polynomials can be initiated from a nonnegative Hermitian matrix-valued measure (or a distribution of bounded variation) on $[0, \infty)$ such that the associated block Hankel matrices $H_{1,j}$ and $H_{2,j}$ are positive definite.

From this distribution, we compute the moments s_j as in (80) and (81). Next, we apply Definition 4. Finally, the second equality in (18) is used to obtain (78).

Next, we recall the characterization of HTM polynomials in terms of block Hankel matrices; see [6, Theorem 7.10].

Theorem 3. *Let n be a natural number with $n \geq 2$, and let $\mathbf{f}_n$ be a matrix-valued polynomial. Let $\mathbf{h}_n$ (resp. $\mathbf{g}_n$) denote the $\mathbf{h}_n$ -part (resp. $\mathbf{g}_n$ -part) of $\mathbf{f}_n$. Furthermore, let $n = 2m$ (resp. $n = 2m + 1$) with $m \geq 1$, and let $H_{1,m-1}$ and $H_{2,m-1}$ (resp. $H_{1,m}$ and $H_{2,m-1}$) be constructed from the Hermitian Markov parameters of the polynomial $\mathbf{f}_n$. Then the matrix-valued polynomial $\mathbf{f}_n$ is an HTM polynomial if and only if the block Hankel matrices $H_{1,m-1}$ and $H_{2,m-1}$ (resp. $H_{1,m}$ and $H_{2,m-1}$) are positive definite.*

We now pass from the Hankel-matrix characterization to a Bezoutian characterization. In particular, for an HTM polynomial, the Bezoutians associated with the decomposition of $\mathbf{f}_n$ into its $\mathbf{h}_n$ - and $\mathbf{g}_n$ -parts have definite signs.

Proposition 3. *Let $\mathbf{f}_n$ be an HTM polynomial. Moreover, let $\mathbf{h}_n$ and $\mathbf{g}_n$ be as in (2), (3) and (4), respectively. Furthermore, let $B_{x\mathbf{g}_n^*, \mathbf{h}_n^*}(x\mathbf{g}_n, \mathbf{h}_n)$ and $B_{\mathbf{g}_n^*, \mathbf{h}_n^*}(\mathbf{g}_n, \mathbf{h}_n)$ be Bezoutians associated with the quadruple $(x\mathbf{g}_n^*, \mathbf{h}_n, \mathbf{h}_n^*, x\mathbf{g}_n)$ and $(\mathbf{g}_n^*, \mathbf{h}_n, \mathbf{h}_n^*, \mathbf{g}_n)$, respectively. Thus,*

- a) *the matrix $B_{x\mathbf{g}_n^*, \mathbf{h}_n^*}(x\mathbf{g}_n, \mathbf{h}_n)$ is positive definite, and*
- b) *the matrix $B_{\mathbf{g}_n^*, \mathbf{h}_n^*}(\mathbf{g}_n, \mathbf{h}_n)$ is negative definite.*

Proof. Let $\tilde{J}_m$, $\tilde{\mathbf{g}}_{2m+1}$, $S(\mathbf{h}_{2m})$, $S(\mathbf{g}_{2m+1})$ and $S(\tilde{\mathbf{g}}_{2m+1})$ be as in (44), (57) and (50). For $n = 2m$, by using (32), (33), (53) and (54), we have

$$\begin{aligned} B_{x\mathbf{g}_{2m}^*, \mathbf{h}_{2m}^*}(x\mathbf{g}_{2m}, \mathbf{h}_{2m}) &= S(\mathbf{h}_{2m}^*)\tilde{J}_{m-1}H_{2,m-1}\tilde{J}_{m-1}S(\mathbf{h}_{2m}), \\ B_{\mathbf{g}_{2m}^*, \mathbf{h}_{2m}^*}(\mathbf{g}_{2m}, \mathbf{h}_{2m}) &= -S(\mathbf{h}_{2m}^*)\tilde{J}_{m-1}H_{1,m-1}\tilde{J}_{m-1}S(\mathbf{h}_{2m}). \end{aligned}$$

By (32), (33), (55) and (56), for $n = 2m + 1$, we obtain

$$\begin{aligned} B_{x\mathbf{g}_{2m+1}^*, \mathbf{h}_{2m+1}^*}(x\mathbf{g}_{2m+1}, \mathbf{h}_{2m+1}) &= S(\mathbf{g}_{2m+1}^*)\tilde{J}_m H_{1,m} \tilde{J}_m S(\mathbf{g}_{2m+1}) \\ B_{\mathbf{g}_{2m+1}^*, \mathbf{h}_{2m+1}^*}(\mathbf{g}_{2m+1}, \mathbf{h}_{2m+1}) &= -S(\tilde{\mathbf{g}}_{2m+1}^*)\tilde{J}_{m-1} H_{2,m-1} \tilde{J}_{m-1} S(\tilde{\mathbf{g}}_{2m+1}). \end{aligned}$$

From Theorem 3, see that assertion a) and b) are valid.

Remark 11. In [40, Theorem 2], it is proven that a scalar polynomial f_n is Hurwitz if and only if $B_{xg_n^*, h_n^*}(xg_n, h_n)$ is negative definite and $B_{g_n^*, h_n^*}(g_n, h_n)$ is positive definite. Note that the Markov parameters related to [40, Theorem 2] have the opposite sign compared with those associated with parts (a) and (b) of Proposition 3.

5. Completing a matrix polynomial to an HTM polynomial

As indicated in [10], not every Hurwitz matrix polynomial is an HTM polynomial. In this section, we propose a procedure to complete a given matrix polynomial into an HTM polynomial.

In [10], the polynomial

$$P_3(z) = I_2 z^3 + A_1 z^2 + A_2 z + A_3 \quad (86)$$

with

$$A_1 = \begin{pmatrix} \frac{2187}{109} & \frac{1206}{109} \\ 0 & 9 \end{pmatrix}, \quad A_2 = \begin{pmatrix} \frac{10662}{109} & \frac{8700}{109} \\ 0 & 18 \end{pmatrix}, \quad A_3 = \begin{pmatrix} \frac{11178}{109} & \frac{10524}{109} \\ 0 & 6 \end{pmatrix} \quad (87)$$

was suggested as a polynomial that is Hurwitz but not an HTM polynomial. The rational matrix function $\frac{1}{z}(A_1 z + A_3)(I_2 z + A_2)^{-1}$ does not satisfy property (30). Moreover, as indicated in [10], the 2×2 matrix coefficients c_j of the expansion

$$\frac{1}{z}(A_1 z + A_3)(I_2 z + A_2)^{-1} = -\frac{c_0}{z} - \frac{c_1}{z^2} - \frac{c_2}{z^3} - \dots$$

are not Hermitian matrices. Consequently, the polynomial P_3 cannot be investigated in the framework of HTM polynomials.

The following result allows one to determine whether a matrix polynomial P_n that is not an HTM polynomial is in fact a Hurwitz polynomial by “completing,” mainly by perturbing P_n with another polynomial Q_{n-1} . This is done by constructing a new HTM polynomial of degree $2n$.

Theorem 4. Let $P_n(z)$ be $q \times q$ a monic polynomial of degree n . If there exists a $q \times q$ polynomial $Q_{n-1}(z)$ of degree $n - 1$ such that

$$\mathbf{f}_{2n}(z) = P_n(z^2) + zQ_{n-1}(z^2) \quad (88)$$

is an HTM polynomial, then the polynomial $P_n(z)$ is a Hurwitz polynomial.

Proof. From (18) and (19), there are polynomials $P_{1,n}$ y $Q_{1,n}$ such that $P_n(z) = (-1)^n P_{1,n}^*(-\bar{z})$ $Q_{n-1}(z) = (-1)^{n+1} Q_{1,n}^*(-\bar{z})$ and $P_{1,n}(z)$ is an orthogonal polynomial on $[0, +\infty)$ with respect to a certain positive measure σ corresponding to the Stieltjes transform $-\frac{Q_{1,n}^*(\bar{z})}{P_{1,n}^*(\bar{z})}$. By [21, Theorem 1], the roots of the polynomial $\det P_{1,n}(z)$ lie in $[0, +\infty)$. Consequently, the roots of the polynomial

$$\det P_n(z) = \det((-1)^n P_{1,n}^*(-\bar{z}))$$

lie in the open left half-plane. Therefore, $\det P_n(z)$ is a Hurwitz polynomial.

Remark 12. Let P_n be a monic matrix polynomial and let $P_n^\top$ denote its transpose. Since

$$\det P_n = \det P_n^\top,$$

Theorem 4 is also valid for $P_n^\top$ and $Q_{n-1}^\top$.

For the construction of the polynomial Q_{n-1} appearing in Theorem 4, we propose the following algorithm. The procedure is based on the representation of the associated rational matrix function in terms of orthogonal polynomials and the corresponding Markov parameters, which allows one to recover Q_{n-1} from a given polynomial P_n .

Algorithm for finding Q_{n-1} from P_n

Step 1. Let $P_{1,n}(z) = (-1)^n P_n^*(-\bar{z})$ (resp. $P_{1,n}(z) = (-1)^n P_n^{\top*}(-\bar{z})$). By using (17), compute $Q_{1,n}(z)$ in terms of the coefficients of $P_n(z)$ (resp. $P_n^\top(z)$) and the moments or Markov parameters s_j , associated with some positive measure σ on $[0, +\infty)$.

Step 2. Verify that $\frac{P_{1,n}^*(\bar{z})}{Q_{1,n}^*(\bar{z})}$ satisfies Equality (30). If Equality (30) is not satisfied, we cannot apply this algorithm.

Step 3. By using the rational matrix function $\frac{P_{1,n}^*(\bar{z})}{Q_{1,n}^*(\bar{z})}$, calculate the remaining Markov parameters from the set $(s_j)_{j=0}^{2n-1}$.

Step 4. If sequence $(s_j)_{j=0}^{2n-1}$ is a positive definite Stieltjes sequence, meaning that the corresponding Hankel matrices $H_{1,n-1}$ and $H_{2,n-1}$ are positive definite matrices, then $Q_{n-1}(z) = (-1)^{n+1} Q_{1,n}^*(-\bar{z})$.

Example 2. We apply the completion procedure to the matrix polynomial (86).

For $P_3^\top(z)$, we use the above algorithm together with Remark 12. With moments $s_0 = \begin{pmatrix} 2b & -b \\ -b & b \end{pmatrix}$, $s_1 = \begin{pmatrix} 4b & -b \\ -b & b \end{pmatrix}$ and $s_2 = \begin{pmatrix} 16b & -2b \\ -2b & 2b \end{pmatrix}$, we calculate $Q_{1,3}(z)$; therefore, we have

$$Q_{1,3}(z) = s_0 z^2 + z(s_1 - A_1^{\top*} s_0) + s_2 - A_1^{\top*} s_1 + A_2^{\top*} s_0.$$

From the Laurent expansion of $-\frac{P_{1,3}^*(\bar{z})}{Q_{1,3}^*(\bar{z})}$, we calculate the moments

$$s_3 = \begin{pmatrix} 96b & -6b \\ -6b & 6b \end{pmatrix}, s_4 = \begin{pmatrix} 768b & -24b \\ -24b & 24b \end{pmatrix}, s_5 = \begin{pmatrix} 7680b & -120b \\ -120b & 120b \end{pmatrix}.$$

The block Hankel matrices $H_{1,2}$ and $H_{2,2}$ are positive definite matrices for $b > 0$. From (19), polynomial $Q_2(z)$ constructed from P_3 has the form $Q_2(z) = Q_{1,3}^*(-\bar{z})$. Thus, the resulting HTM polynomial $\mathbf{f}_6$ can be represented as in (88):

$$\mathbf{f}_6(z) = z^6 I_2 + B_1 z^5 + A_1 z^4 + B_2 z^4 + A_2 z^2 + B_3 z + A_3. \quad (89)$$

The coefficients A_1 , A_2 and A_3 are given by (87), and the coefficients B_1 , B_2 and B_3 are

$$B_1 = \begin{pmatrix} 2b & -b \\ -b & b \end{pmatrix}, B_2 = \begin{pmatrix} \frac{2732b}{109} & -8b \\ -8b & 8b \end{pmatrix}, B_3 = \begin{pmatrix} \frac{6826b}{109} & -11b \\ -11b & 11b \end{pmatrix}.$$

By Theorem 2, matrix polynomial (89) is a Hurwitz matrix polynomial for $b > 0$. Finally, by Theorem 4, polynomial P_3 as in (86) is a Hurwitz matrix polynomial.

In the scalar version, one can construct a sequence of scalar Hurwitz polynomials $(f_j)_{j=1}^\infty$ using the sequence of scalar orthogonal polynomials on $[0, +\infty)$ $(p_{1,j})_{j=1}^\infty$ and $(p_{2,j})_{j=1}^\infty$ as in Definition 4; see [12].

In the following lemma, we propose constructing a family of scalar Hurwitz polynomials involving the polynomials $p_{1,j}$, and $p_{2,j}$ and their derivatives.

Lemma 4. For $q = 1$, let $(p_{1,m}(z))$ (resp. $(p_{2,m}(z))$) be as in Definition 4, and let $\tilde{p}_{1,m-1}(z) := \frac{d}{dz} p_{1,m}(z)$ (resp. $\tilde{p}_{2,m-1}(z) := \frac{d}{dz} p_{2,m}(z)$). Thus, for $m \geq 2$, the polynomials

$$f_{2m}(z) = (-1)^m (p_{1,m}(-z^2) - z \tilde{p}_{1,m-1}(-z^2)) \quad (90)$$

and

$$f_{2m+1}(z) = (-1)^m (\tilde{p}_{2,m}(-z^2) + z p_{2,m}(-z^2)). \quad (91)$$

are scalar Hurwitz polynomials.

Proof. The proof of (90) and (91) readily follows from the interlacing properties of the roots of the orthogonal polynomials $p_{k,m}(z)$ and $\tilde{p}_{k,m-1}(z)$ for $k = 1, 2$, and from the fact that $a_0 a_1 > 0$, where a_0 and a_1 are the leading coefficients of the polynomial $f_n(z) = a_0 z^n + a_1 z^{n-1} + \dots + a_n$; see the Hermite–Biehler theorem [3], [32].

Remark 13. From (78) and (89), we see that the determinants of the coefficients of these polynomials are positive. Based on this observation, we propose the conjecture below.

Conjecture. The determinants of the coefficients A_j of an HTM polynomial $\mathbf{f}_n$ are positive.

6. Additional remarks

In this section, to clarify the importance of the present work, we recall several relevant results from [49], paying particular attention to certain steps in the proofs that were left implicit. We emphasize that the authors of [49] (2021) were the first

to propose that the HTM polynomials possess the Hurwitz property, while the concept of HTM polynomials itself was introduced earlier by the author of the present work in [6] (2015).

Lemma 5. [49, Lemma 3.4] *Let $N_R(z)$, $D_R(z)$, $N_L(z)$ and $D_L(z) \in \mathbb{C}[z]^{p \times p}$ be regular such that $\deg D_R > \deg N_R$, $\deg D_L > \deg N_L$ and, for all large enough $z \in \mathbb{C}$,*

$$(D_L(z))^{-1}N_L(z) = N_R(z)(D_R(z))^{-1} = \sum_{k=0}^{\infty} z^{-(k+1)} s_k.$$

If $D_L(z)$ and $D_R(z)$ are written in the form

$$D_L(z) = \sum_{k=0}^{m_L} D_{L,m_L-k} z^k \quad \text{and} \quad D_R(z) = \sum_{k=0}^{m_R} D_{R,m_R-k} z^k,$$

where $D_{L,k} \in \mathbb{C}^{p \times p}$ for $k = 0, \dots, m_L$ and $D_{R,k} \in \mathbb{C}^{p \times p}$ for $k = 0, \dots, m_R$, then

$$\mathbf{B}_{D_L, N_L}(D_R, N_R) = \begin{pmatrix} D_{L,m_L-1} & \cdots & D_{L,0} \\ \vdots & \ddots & \vdots \\ D_{L,0} & \cdots & \end{pmatrix} \begin{pmatrix} s_0 & \cdots & s_{m_R-1} \\ \vdots & \ddots & \vdots \\ s_{m_L-1} & \cdots & s_{m_R+m_L-2} \end{pmatrix} \cdot \begin{pmatrix} D_{R,m_R-1} & \cdots & D_{R,0} \\ \vdots & \ddots & \vdots \\ D_{R,0} & \cdots & \end{pmatrix}. \quad (92)$$

Let L_1 and L be as in (74). In [49], the Theorem 3.5 is regarding the inertia representation of matrix polynomials $\mathbf{f}_n$. For $n = 2m$, in the proof of [49, Theorem 3.5] the [49, Equality (3.6)] in terms of our notations is the following

$$B_{L_1^*, L^*}(L, L_1) = 2i B_{\mathbf{h}_{2m}^*(-\bar{x}^2), x\mathbf{g}_{2m}^*(-\bar{x}^2)}(\mathbf{h}_{2m}(-x^2), -x\mathbf{g}_{2m}(-x^2)). \quad (93)$$

The authors of [49] use $2i B_{F_e^\vee, -F_0^\vee}(\widehat{F}_e, -\widehat{F}_0)$ in place of the right-hand side of (93). They claim that Lemma 5 implies that the matrix $2i B_{F_e^\vee, -F_0^\vee}(\widehat{F}_e, -\widehat{F}_0)$ is congruent to the block Hankel matrix

$$\begin{pmatrix} s_0 & 0_p & s_1 & \cdots & s_{m-1} & 0_p \\ 0_p & s_1 & 0_p & \cdots & 0_p & s_m \\ s_1 & 0_p & s_2 & \cdots & s_m & 0_p \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ s_{m-1} & 0_p & s_m & \cdots & s_{2m-2} & 0_p \\ 0_p & s_m & 0_p & \cdots & 0_p & s_{2m-1} \end{pmatrix}, \quad (94)$$

and that, after a suitable reordering of rows and columns in (94), it is further congruent to

$$\begin{pmatrix} H_{1,m-1} & 0 \\ 0 & H_{2,m-1} \end{pmatrix}. \quad (95)$$

Remark 14. *By comparing the Hankel matrix of the right-hand side of (92) and the matrix (94) It is not clear how from $2i B_{F_e^\vee, -F_0^\vee}(\widehat{F}_e, -\widehat{F}_0)$ one gets (94). Furthermore, it is not clear how from (94) one obtains (95).*

In contrast, for $n = 2m$, in the present work we derive the matrix $\begin{pmatrix} H_{2,m-1} & 0 \\ 0 & H_{1,m-1} \end{pmatrix}$ associated with the quadratic form in (95) by employing the decomposition (34) and by computing explicitly the quadratic forms corresponding to (32) and (33), as shown in (66). Additionally, for $n = 2m + 1$, the proof of [49, Theorem 3.5] is not given.

In Theorem 4.10 of [49], a “stability criterion via Markov parameters” for matrix polynomials $\mathbf{f}_n$ in even and odd cases is suggested. We have the following remark regarding this.

Remark 15. *For $n = 2m$, Theorem 4.10 of [49], is related to the HTM polynomials. Other matrix polynomials, for example, the Hurwitz polynomial (86), are not covered by [49, Theorem 4.10].*

In this work, we do not discuss the case $n = 2m + 1$ of Theorem 4.10 of [49].

7. Conclusion

In this work we obtained an explicit representation of the Bezoutian associated with Hurwitz-type matrix polynomials. This representation clarifies the relation between Hurwitz-type matrix polynomials and classical Hurwitz matrix polynomials and provides a constructive tool for verifying Hurwitzness in the matricial setting. Our approach, based on the decomposition of Bezoutians and on the structure of truncated matricial Stieltjes moment problems, offers a natural framework for extending classical stability criteria to matrix polynomials. In addition, we proposed a method to enlarge the class of Hurwitz-type matrix polynomials by perturbing a polynomial that is not of Hurwitz type so that the new polynomial becomes Hurwitz type. These results contribute to a better understanding of the structure of Hurwitz-type matrix polynomials and their relation to Bezoutians, and they suggest further developments in matrix moment problems, continued fractions, and stability theory.

Acknowledgments

This work was partially supported by the Universidad Michoacana de San Nicolás de Hidalgo and SNII–SECIHTI grants, México.

Conflicts of Interest: The authors declare no conflict of interest.

REFERENCES

1. N. I. Aheizer, M. G. Krein, Some Questions in the Theory of Moments. Translations of Mathematical Monographs, 2, American Mathematical Society, Providence, R.I., 1962, 265 pp. <https://doi.org/10.1090/mmono/002>

2. B. D. O. Anderson, E. I. Jury, Generalized Bezoutin and Sylvester matrices in multivariable linear control, *IEEE Trans. Autom. Control.* – 1976. – Vol. **AC-21**. – P. 551–556. <https://doi.org/10.1109/TAC.1976.1101263>
3. C. Biehler, Sur une classe d'équations algébriques dont toutes les racines sont réelles, *J. Reine Angew. Math. (Crelle's Journal)* – 1879. – Vol. **87**. – P. 350–352. <https://doi.org/10.1515/9783112341889-020>
4. D. Cox, J. Little, and D. O'Shea, *Ideals, Varieties, and Algorithms*, 4th ed., *Undergraduate Texts in Mathematics*, Springer, Cham. – 2015. <https://doi.org/10.1007/978-3-031-91841-4>
5. A.E. Choque Rivero, On Dyukarev's resolvent matrix for a truncated Stieltjes matrix moment problem under the view of orthogonal matrix polynomials, *Linear Algebra Appl.* – 2015. – Vol. **474**. – P. 44–109. <https://doi.org/10.1016/j.laa.2015.01.027>
6. A.E. Choque Rivero, On matrix Hurwitz type polynomials and their interrelations to Stieltjes positive definite sequences and orthogonal matrix polynomials, *Linear Algebra Appl.* – 2015. Vol. **476**. – P. 56–84. <https://doi.org/10.1016/j.laa.2015.03.001>
7. A.E. Choque-Rivero, The Kharitonov theorem and robust stabilization via orthogonal polynomials, *Visnyk of V. N. Karazin Kharkiv National University. Ser. Mathematics, Applied Mathematics and Mechanics.* – 2017. Vol. **86**. – P. 49–68. <https://doi.org/10.26565/2221-5646-2017-86-05>
8. A.E. Choque Rivero, Hurwitz polynomials and orthogonal polynomials generated by Routh–Markov parameters, *Mediterr. J. Math.* – 2018. Vol. **15**, No **40**. – 15 pages. <https://doi.org/10.1007/s00009-018-1083-2>
9. A.E. Choque Rivero, Extended set of solutions of a bounded finite-time stabilization problem via the controllability function, *IMA J. Math. Control Inf.* – 2021. Vol. **38**, No **4**. – P. 1174–1188. <https://doi.org/10.1093/imamci/dnab028>
10. A.E. Choque Rivero, Comments on the paper “On the relation between Hurwitz stability of matrix polynomials and matrix-valued Stieltjes functions,” *J. Comput. Appl. Math.* 417 (2023) by Xuzhou Zhan and Yongjian Hu, submitted to *J. Comput. Appl. Math.*, 2023. Available at: https://drive.google.com/file/d/1qE3xsF37K9HN2lrMQ9eW92pH9LEk_6Te/view?usp=sharing
11. A.E. Choque Rivero, Korobov's controllability function as motion time: Extension of the solution set of the synthesis problem, *Zh. Mat. Fiz. Anal. Geom.* – 2023. Vol. **19**, No **3**. – P. 556–586. <https://doi.org/10.15407/mag19.03.556>

12. A.E. Choque-Rivero, I. Area, Favard type theorem for Hurwitz polynomials, *Discrete & Continuous Dynamical Systems–B.* – 2020. Vol. **22**, No **2**. – P. 529–544. <https://doi.org/10.3934/dcdsb.2019252>
13. A.E. Choque Rivero, V.I. Korobov, V.O. Skoryk, Controllability function as time of motion. I, *Mat. Fiz. Anal. Geom.* – 2004. Vol. **2**, No **11**. – P. 208–225. (in Russian) Translated into English in <http://arxiv.org/abs/1509.05127>.
14. A.E. Choque Rivero, V.I. Korobov, V.O. Skoryk, Controllability function as time of motion. II, *Mat. Fiz. Anal. Geom.* – 2004. Vol. **3**, No **11**. – P. 341–354.
15. A.E. Choque Rivero, C. Maedler, On resolvent matrix, Dyukarev–Stieltjes parameters and orthogonal matrix polynomials via $[0, +\infty)$ –Stieltjes transformed sequences, *Complex Anal. Oper. Theory.* – 2019. Vol. **13**. – P. 1–44. <https://doi.org/10.1007/s11785-017-0655-7>
16. B.N. Datta, Application of Hankel matrices to Markov parameters to the solution of the Routh–Hurwitz and the Schur–Cohn problems, *J. Math. Anal. Appl.* – 1979. Vol. **68**, No **1**. – P. 276–290. [https://doi.org/10.1016/0022-247x\(79\)90115-x](https://doi.org/10.1016/0022-247x(79)90115-x)
17. J.E. Dennis, J.F. Traub and R.P. Weber, On the Matrix Polynomial, Lambda-Matrix and Block Eigenvalue Problems, Computer Science Department Tech. rep., Cornell Univ., Ithaca, N.Y., and Carnegie-Mellon Univ., Pittsburgh, Pa. – 1971. <https://hdl.handle.net/1813/5954>
18. Y.M. Dyukarev, A general scheme for solving interpolation problems in the Stieltjes class that is based on consistent representations of pairs of nonnegative operators. I. *Mat. Fiz. Anal. Geom.* – 1999. Vol **6**, No **1–2**. – P. 30–54. <https://mag.ilt.kharkiv.ua/index.php/mag/article/view/m06-0030r>
19. Y.M. Dyukarev, Indeterminacy criteria for the Stieltjes matrix moment problem, *Math Notes.* – 2004. Vol **75**. – P. 66–82. <https://doi.org/10.1023/B:MATN.0000015022.02925.bd>
20. Y.M. Dyukarev, Theory of Interpolation Problems in the Stieltjes Class and Related Problems of Analysis, Habilitation thesis. Kharkiv National University – 2006.
21. Y.M. Dyukarev, The zeros of determinants of matrix-valued polynomials that are orthonormal on a semi-infinite or finite interval, *Sbornik: Mathematics.* – 2018. Vol. **209**, No **12**. – P. 1745–1755.
22. Y.M. Dyukarev, B. Fritzsche, B. Kirstein, C. Mädler, On truncated matricial Stieltjes type moment problems. *Complex Anal. Oper. Theory.* – 2010. Vol. **4**. – P. 905–951. <https://doi.org/10.1007/s11785-009-0002-8>

23. Y.M. Dyukarev, B. Fritzsche, B. Kirstein, C. Mädler, H. Thiele, On distinguished solutions of truncated matricial Hamburger moment problems. *Complex Anal. Oper. Theory.* – 2009. Vol. **3**, No. **759**. – P. 759–834. <https://doi.org/10.1007/s11785-008-0061-2>
24. B. Fritzsche, B. Kirstein, and Conrad Mädler, On Hankel nonnegative definite sequences, the canonical Hankel parametrization, and orthogonal matrix polynomials, *Complex Anal. Oper. Theory.* – 2011. Vol. **5**, No. **2**. – P. 447–511. <https://doi.org/10.1007/s11785-010-0054-9>
25. B. Fritzsche, B. Kirstein, and C. Madler, On a special parametrization of matricial α –Stieltjes one-sided non-negative definite sequences. *Interpolation, Schur functions and moment problems. II. Oper. Theory Adv. Appl.* – 2012. Vol. **226**. – P. 211–250. https://doi.org/10.1007/978-3-0348-0428-8_5
26. B. Fritzsche, B. Kirstein, and Conrad Mädler, Transformations of matricial α –Stieltjes non-negative definite sequences, *Linear Algebra Appl.* – 2013. Vol. **439**, No. **12**. – P. 3893–3933. <https://doi.org/10.1016/j.laa.2013.10.002>
27. P. A. Fuhrmann, B. N. Datta, On Bezoutians, Van der Monde matrices and the Lienard–Chipart stability criterion, *Linear Algebra Appl.* – 1989. Vol. **120**. – P. 23–37. [https://doi.org/10.1016/0024-3795\(89\)90367-4](https://doi.org/10.1016/0024-3795(89)90367-4)
28. F.R. Gantmacher, *The Theory of Matrices*, Volume I, AMS Chelsea. – 2000.
29. F.R. Gantmacher, *The Theory of Matrices*, Volume II, AMS Chelsea. – 2000.
30. G. Heinig, U. Jungnickel, On the Routh–Hurwitz and Shur–Cohn problems for matrix polynomials and generalized Bezoutians, *Math. Nach.* – 1984. Vol. **116**, No. **1**. – P. 185–196. <https://doi.org/10.1002/mana.19841160114>
31. D. Henrion, D. Arzelier, D. Peaucelle, M. Sebek, An LMI condition for robust stability of polynomial matrix polytopes, *Automatica.* – 2001. Vol. **37**, No. **3**. – P. 461–468. [https://doi.org/10.1016/S0005-1098\(00\)00170-9](https://doi.org/10.1016/S0005-1098(00)00170-9)
32. C. Hermite, Extrait d’une lettre de. M. Hermite de Paris à M. Borchardt de Berlin sur le nombre des racines d’une équation algébrique comprises entre des limites données, *J. Reine Angew. Math.* – 1856. Vol. **1856**, No. **52**. – P. 39–51. <https://doi.org/10.1515/crll.1856.52.39>
33. O. Holtz, M. Tyaglov, Structured matrices, continued fractions, and root localization of polynomials. *SIAM Rev.* – 2012. Vol. **54**, No. **3**. – P. 421–509. <https://doi.org/10.1137/090781127>
34. A. Hurwitz, Über die Bedingungen unter welchen eine Gleichung nur Wurzeln mit negativen reellen Teilen besitzt *Math. Ann.* – 1895. Vol. **46**. – P. 273–284. <https://doi.org/10.1007/BF01446812>

35. E.J. Jury, A. Katbab, A note on Kharitonov-type results in the space of Markov parameters, *IEEE Trans. Autom. Control.* – 1989. Vol. **37**, No **1**. – P. 155–158. <https://doi.org/10.1109/9.109655>
36. V. Katsnelson, Stieltjes functions and Hurwitz stable entire functions, *Complex Anal. Oper. Theory.* – 2011. Vol. **5**. – P. 611–630. <https://doi.org/10.1007/s11785-011-0146-1>
37. F.J. Kraus, M. Mansour and M. Sebek, Hurwitz matrix for polynomial matrices, In: Jeltsch, R., Mansour, M. (eds) *Stability Theory. ISNM International Series of Numerical Mathematics*, Birkhäuser Basel. – 1996. Vol. **121**. – P. 67–74. https://doi.org/10.1007/978-3-0348-9208-7_8
38. M.G. Krein, M.A. Naimark, The method of symmetric and Hermitian forms in the theory of the separation of the roots of algebraic equations, *Linear and Multilinear Algebra.* – 1981. Vol. **10**, No **4**. – P. 265–308. <https://doi.org/10.1080/03081088108817420>
39. P. Lancaster, *Lambda-Matrices and Vibrating Systems*, Pergamon Press, New York. – 1966. <https://doi.org/10.1016/C2013-0-01959-7>
40. P. Lancaster and M. Tismenetsky, *The Theory of Matrices: With Applications*, Second Edition, Academic Press. – 1985.
41. L. Lerer, M. Tismenetsky, The Bezoutian and the eigenvalue-separation problem for matrix polynomials, *Integral Equations Oper. Theory.* – 1982. Vol. **5**. – P. 386–445. <https://doi.org/10.1007/BF01694045>
42. L. Lerer, L. Rodman, M. Tismenetsky, Inertia theorems for matrix polynomials, *Linear and Multilinear Algebra.* – 1991. Vol. **30**, No **3**. P. 157–182. <https://doi.org/10.1080/03081089108818100>
43. C. MacDuffee, *The Theory of Matrices*, Chelsea, New Your. – 1946.
44. F. Martins, E. Pereira, Block matrices and stability theory, *Tatra Mt. Math. Publ.* – 2007. Vol. **38**. – P. 147–162.
45. D.V. Ouellette, Schur complements and statistics, *Linear Algebra Appl.* – 1981. Vol. **36**. – P. 187–295. [https://doi.org/10.1016/0024-3795\(81\)90232-9](https://doi.org/10.1016/0024-3795(81)90232-9)
46. P. Resende and E. Kaszkurewicz, New stability conditions for matrix polynomials using a block-bidiagonal form, *IEEE International Symposium on Circuits and Systems*, Espoo, Finland. – 1988. Vol. **1**. – P. 385–388. <https://doi.org/10.1109/iscas.1988.14945>
47. E. J. Routh, *A Treatise on the Stability of a Given State of Motion: Particularly Steady Motion*. Macmillan and Co. – 1877.

48. L. Shieh, S. Sacheti, A matrix in the block Schwarz form and the stability of matrix polynomials, *Int. J. Control.* – 1978. Vol. **27**, No **2**. – P. 245–259. <https://doi.org/10.1080/00207177808922362>
49. X. Zhan, A. Dyachenko, On generalization of classical Hurwitz stability criteria for matrix polynomials, *J. Comput. Appl. Math.* – 2021. Vol. **383**. – P. 1–16. <https://doi.org/10.1016/j.cam.2020.113113>

Article history: Received: 27 February 2026; Final form: 10 May 2026

Accepted: 13 May 2026. Published 31 May 2026.

How to cite this article:

A. E. Choque–Rivero, On the Hurwitz stability of Hurwitz-type matrix polynomials, *Visnyk of V. N. Karazin Kharkiv National University. Ser. Mathematics, Applied Mathematics and Mechanics*, Vol. 103, 2026, p. 5–37. <https://doi.org/10.26565/2221-5646-2026-103-01>

Про стійкість за Гурвіцем матричних поліномів типу Гурвіца

А. Е. Чоке-Рівер

Інститут фізики та математики

Університет Мічоакана-де-Сан-Ніколас-де-Ідальго

Будівля С-3, Центральний кампус, 58060, Морелія, Мічоакан, Мексика

Дійсний скалярний поліном, корені якого лежать у лівій півплощині комплексної площини, називається поліномом Гурвіца. Це поняття походить з робіт Дж. К. Максвелла, Е. Дж. Раута, та А. Гурвіца, і пізніше вивчалось за допомогою таких методів, як послідовності Штурма, параметри Маркова, та ланцюгові дробі; див., наприклад, *Теорія матриць*, Том II, Розділ XV, AMS Chelsea (2000), Ф. Гантмахера.

Матричний поліном $P(z)$ розміру $q \times q$ називається Гурвіцевим якщо $\det P(z)$ є Гурвіцевим поліномом. Кожен матричний поліном $\mathbf{f}_n$ може бути представлений у формі $\mathbf{f}_n(z) = \mathbf{h}_n(z^2) + z\mathbf{g}_n(z^2)$. Матричний поліном $\mathbf{f}_{2m}$ називається поліномом типу Гурвіца якщо вираз $\mathbf{g}_{2m}(z)\mathbf{h}_{2m}^{-1}(z)$ допускає представлення у вигляді скінченного ланцюгового дробу з додатно визначеними матричними коефіцієнтами. Аналогічно, матричний поліном непарного степеня $\mathbf{f}_{2m+1}$ є поліномом типу Гурвіца якщо $\frac{1}{z}\mathbf{h}_{2m+1}(z)\mathbf{g}_{2m+1}^{-1}(z)$ має таку ж саму властивість. Поняття матричних поліномів типу Гурвіца було введено в *Про матричні поліноми типу Гурвіца та їх взаємозв'язки з додатно визначеними послідовностями Стилтьєса та ортогональними матричними поліномами*, *Linear Algebra Appl.* 476 (2015), автором А. Е. Чоке-Ріверо.

У цій роботі ми виводимо явну форму Безутіана, пов'язаного з матричними поліномами типу Гурвіца. Той факт, що матричні поліноми типу Гурвіца є матричними поліномами Гурвіца, був запропонований та частково доведений з використанням Безутіанів у *Про узагальнення класичних критеріїв стійкості Гурвіца для матричних поліномів*, *J. Comput. Appl. Math.* 383 (2021), авторами Х. Жаном та А. Дьяченком. На відміну від цієї роботи, ми використовуємо розкладання форми Безутіана, введене в *Деякі питання теорії моментів*, Переклади математичних

монографій 2, AMS, 1962, авторами Н. І. Ахієзер та М. Г. Крейн, для скалярних поліномів.

Крім того, ми пропонуємо метод розширення класу матричних поліномів типу Гурвіца шляхом додавання до заданого полінома матричного полінома, який не є поліномом типу Гурвіца, але при цьому отриманий поліном стає типу Гурвіца.

Ключові слова: Гурвіцевий матричний поліном; ортогональний матричний поліном; Безутіан; матричний поліном типу Гурвіца.

Конфлікт інтересів: Автори повідомляють про відсутність конфлікту інтересів.

Історія статті: отримана: 27 лютого 2026; останній варіант: 10 травня 2026
прийнята: 13 травня 2026. Оприлюднено 31 травня 2026.